

EGU25-2716, updated on 11 Apr 2026

<https://doi.org/10.5194/egusphere-egu25-2716>

EGU General Assembly 2025

© Author(s) 2026. This work is distributed under the Creative Commons Attribution 4.0 License.



## Evaporation dynamics from flowing water surfaces

Lintong Hou<sup>1</sup>, Milad Aminzadeh<sup>1,2</sup>, Dani Or<sup>3,4</sup>, Justus Patzke<sup>5</sup>, Peter Fröhle<sup>5</sup>, and Nima Shokri<sup>1,2</sup>

<sup>1</sup>Institute of Geo-Hydroinformatics, Hamburg University of Technology, Hamburg, Germany

<sup>2</sup>United Nations University Hub on Engineering to Face Climate Change at the Hamburg University of Technology, United Nations University Institute for Water, Environment and Health (UNU-INWEH), Hamburg, Germany

<sup>3</sup>Soil and Terrestrial Environmental Physics, ETH Zurich, Zurich, Switzerland

<sup>4</sup>Department of Civil and Environmental Engineering, University of Nevada, Reno, NV, USA

<sup>5</sup>Institute of River and Coastal Engineering, Hamburg University of Technology, Hamburg, Germany

In contrast to the wealth of information on evaporation dynamics from placid water surfaces such as lakes and reservoirs, estimating water evaporation from turbulent surfaces of streams remains a challenge. Evidence suggests a considerable change in evaporation from flowing surfaces relative to placid surfaces with local modifiers such as chemical, physical and biological processes that alter the energy budget and water temperature. While the studies on evaporation from wavy surfaces of oceans offer valuable insights, significant differences in hydrodynamics and heat exchange processes distinguish evaporation in oceans from that in rivers. Here we experimentally investigate how water flow characteristics (velocity and turbulence) and atmospheric boundary conditions (wind and radiation) affect evaporation rates and temperature dynamics in a flume. A closed flume (7.6 m length, 0.31 m width, and 0.5 m depth) is used to impose different boundary conditions over a test section of the flume (length of 1.5 m) while other parts of the flume are covered to reduce evaporative losses. Our preliminary findings show significant enhancement in evaporation rates, reaching 2-5 times that of placid water surfaces, driven by increases in surface velocity and turbulence characteristics. Furthermore, we observe that radiative and aerodynamic factors contribute nonlinearly to evaporation enhancement and affect temperature distribution in the water body. The study offers novel insights into evaporation from wavy and turbulent flowing water surfaces for better prediction of evaporation from riverine networks across flow regimes and climatic conditions.