

Mineralogical control of Se and Te signatures in peridotites: implications for the primitive mantle

S. König (1), J.-P. Lorand (2), A. Luguet (1), F. Wombacher (3), and D.G. Pearson (4)

(1) Universität Bonn, Steinmann Institut, Germany (stephan.koenig@uni-bonn.de), (2) Laboratoire de Planétologie et Géodynamique, University de Nantes, France, (3) Universität zu Köln, Institut für Geologie und Mineralogie, Germany, (4) University of Alberta, Department of Earth and Atmospheric Sciences, Edmonton, Canada

Selenium and tellurium belong to the group of highly siderophile elements (HSE) that are believed to constitute key tracers for planetary processes such as formation of the Earth's core and the Late Veneer composition. Constraints on HSE systematics on the planetary scale require, however, a previous understanding regarding the behaviour of these elements during petrogenetic processes and their abundances in the primitive upper mantle (PUM). Recent studies have shown systematic differences in Se/Te between fertile lherzolites and depleted harzburgites. In contrast to fertile lherzolites which remain at broadly chondritic values [1], depleted peridotites are highly fractionated with up to suprachondritic Se/Te (up to 35), where high Se/Te correlates with decreasing Te concentrations [2]. Fractionation is also observed at the scale of a single sample and results from the heterogeneous distribution of micrometer sized Te-bearing host phases in residual peridotites. On the whole rock scale this effect results in a more incompatible behaviour of Te compared to Se, once base metal sulfides are highly depleted and in some cases entirely consumed by partial melting. The marked differences in Se-Te systematics observed between fertile lherzolites and depleted harzburgites can be explained by the combined effect of i) different abundances and proportions of residual and metasomatic base metal sulfides and ii) discrete micrometric to nanometric platinum-group minerals. Hence, it is critical to fully understand Se-Te systematics in harzburgites if we are to provide new insights into the behaviour of Se and Te during mantle depletion. This knowledge is a prerequisite to further constrain the Se and Te abundances of the primitive upper mantle.

[1] Lorand and Alard (2010) Chem Geol 278, 120-130. [2] König et al. (2012) GCA, 86 354–366.