



Impact of radar data assimilation on WRF simulations of the Aniene flood

I. Maiello (1,2), R. Ferretti (2), S. Gentile (2), M. Montopoli (3), E. Picciotti (4), G. Giuliani (2), and F.S. Marzano (5)

(1) Department of Engineering and Environmental Physics, University of Basilicata, (delfida2003@yahoo.it), (2) Centre of Excellence CETEMPS, Department of Physics - Atmospheric Sciences University of L'Aquila, (3) University of L'Aquila/CETEMPS Department of Electrical and Information Engineering, (4) HIMET s.r.l., (5) CETEMPS/Department of Electronic Engineering, Sapienza University of Rome

A study of the Aniene flood, in the urban area of Rome, is performed using Monte Midia Radar data to improve high resolution initial conditions. The radar site is at the border between the Abruzzo and Lazio regions in Central Italy. Mt. Midia top height is at 1760 m and covering most Central Italy, including the urban area of Rome. Aniene flood occurred during May 20-22, 2008 causing severe damages. The meteorological structure was characterized by a deep cyclone impinging on the Tyrrhenian sea for almost two days. During the event radar detected either local convection northern of Rome and moderate rain east of Rome lasting for several hours. High resolution simulations using WRFV3 model are performed; to the aim of improving the Initial Conditions radar data have been assimilated using 3DVAR. Sensitivity tests to different set of Initial Conditions are performed using either ECMWF analyses and 'warm start'. Moreover, 'ad hoc' radar reflectivity coefficients for the Monte Midia radar have been used to analyze their impact in the model forecast. Finally, to objectively identify the best IC statistical indicators are used as FBias, RMS and EQTS for the accumulated precipitation.