



## Climatological evaluation of the COST733 database of circulation classifications using multiple criteria

R. Huth (1,2), C. Beck (3), and M. Cahynová (1)

(1) Institute of Atmospheric Physics, Dept. of Climatology, Prague, Czech Republic (huth@ufa.cas.cz, +420 2 72763745), (2) Charles University, Faculty of Science, Dept. of Physical Geography and Geoecology, Prague, Czech Republic, (3) University of Augsburg, Institute of Geography, Augsburg, Germany

The database of classifications of atmospheric circulation patterns, produced within the COST733 Action, consists in its version v2.0 of 423 different classifications for each of 12 domains over Europe. The classifications differ from each other in five characteristics: (i) the classification method used, (ii) in the number of types, (iii) the classified variable, (iv) sequentiality (whether instantaneous circulation patterns or their 4-day sequences are classified), and (v) the seasonality of definition (whether classifications are defined for a whole year or separately for each season). The goal of the study is to evaluate how successful are the classifications in characterizing (stratifying) surface climate elements, viz., daily minimum and maximum temperature and precipitation. To this end, we employ several criteria: explained variance, pseudo-F statistic, Kolmogorov-Smirnov statistic, confidence interval of the mean, and Silhouette index. Both station and gridded datasets are used. The classifications are ranked in each domain separately. This allows one to evaluate the effect the options (i) to (v) have on the quality of classifications in terms of the stratification of surface temperature and precipitation, to compare the performance among the criteria, and to assess the geographical dependence of the findings.