EMS Annual Meeting Abstracts Vol. 11, EMS2014-509-1, 2014 14th EMS / 10th ECAC © Author(s) 2014



## Comparison of Energy- and Flux-Budget turbulence closure with Large Eddy Simulation

Evgeny Kadantsev (1,2), Carl Fortelius (2), Andrey Glazunov (3), Sergej Zilitinkevich (1,2,4) (1) University of Helsinki, Finland, (2) Finnish Meteorological Institute, Finland, (3) Institute of Numerical Mathematics RAS, Russia, (4) Nizhny Novgorod State University, Russia

Recently a new theoretical derivation of for a new turbulence closure has been presented by Zilitinkevich et al. (2013). This Energy- and Flux-Budget (EFB) closure is based on the budget equations for the basic second moments: the two energies, the TKE  $E_K$  and the turbulent potential energy (TPE)  $E_P$ , and the vertical turbulent fluxes of momentum and potential temperature,  $\tau_i$  (i=1,2) and  $F_z$ . Instead of the traditional postulation of the down-gradient turbulent transport, it uses the flux-profile relationships and determines the eddy viscosity and eddy conductivity from the steady-state version of the budget equations for  $\tau_i$  and  $F_z$ . In that closure authors further advance the physical background of the EFB closure, introduce a new prognostic equation for the turbulent dissipation time scale  $t_T$ , and extend the theory to non-steady turbulence regimes accounting for non-gradient and non-local turbulent transports (when the traditional concepts of eddy viscosity and eddy conductivity become generally inconsistent).

In the present study we implement a new turbulence closure in one-column RANS model and in HAR-MONIE/AROME weather prediction system (HIRLAM-B, 2013). We test the closure in various idealized cases, varying stratification from stable (GABLS1 case) to neutral (Truly Neutral and Conventionally Neutral cases) running HARMONIE/AROME model in single-column mode. Results are compared with LES runs and different numerical weather prediction models.

Bougeault P., and P. Lacarrere, 1989: Parameterization of orography-induced turbulence in a meso-beta scale model. Mon. Wea. Rev., 117, 1872-1890.

Cuxart, J., P. Bougeault, and J.-L. Redelsperger, 2000: A turbulence scheme allowing for mesoscale and large-eddy simulations. Quart. J. Roy. Meteor. Soc., 126, 1-30.

HIRLAM-B Project, 2013: General description of the HARMONIE model. http://hirlam.org/index.php?option=com\_content&view=article&id=65&Itemid=102

Zilitinkevich S.S., T. Elperin, N. Kleeorin, I. Rogachevskii, and I. Esau, 2013: A Hierarchy of Energy- and Flux-Budget (EFB) Turbulence Closure Models for Stably-Stratified Geophysical Flows. Boundary-Layer Meteorol, 146, 341-373.