



EMS Annual Meeting Abstracts

Vol. 19, EMS2022-701, 2022

<https://doi.org/10.5194/ems2022-701>

EMS Annual Meeting 2022

© Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.



## Gaussian-process emulation for integrating data-driven aerosol-cloud physics from simulation, satellite, and ground-based data

Franziska Glassmeier<sup>1</sup>, Fabian Hoffmann<sup>2,3</sup>, Graham Feingold<sup>4</sup>, Edward Gryspeerdt<sup>5</sup>, Antoon van Hooft<sup>1</sup>, Takano Yamaguchi<sup>3,4</sup>, Jill S. Johnson<sup>6</sup>, and Ken S. Carslaw<sup>6</sup>

<sup>1</sup>Department Geoscience and Remote Sensing, TU Delft, Delft, Netherlands

<sup>2</sup>Meteorologisches Institut, Ludwig-Maximilians-Universität München, Munich, Germany

<sup>3</sup>Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder (CO), USA

<sup>4</sup>Chemical Sciences Laboratory, NOAA/Earth System Research Laboratories, Boulder (CO), USA

<sup>5</sup>Space and Atmospheric Physics Group, Imperial College London, London, UK

<sup>6</sup>School of Earth and Environment, University of Leeds, Leeds, UK

Data-driven quantification and parameterization of cloud physics in general, and of aerosol-cloud interactions in particular, rely on input data from observations or detailed simulations. These data sources have complementary limitations in terms of their spatial and temporal coverage and resolution; simulation data has the advantage of readily providing causality but cannot represent the full process complexity. In order to base data-driven approaches on comprehensive information, we therefore need ways to integrate different data sources.

We discuss how the classical statistical technique of Gaussian-process emulation can be combined with specifically initialized ensembles of detailed cloud simulations (large-eddy simulations, LES) to provide a framework for evaluating data-driven descriptions of cloud characteristics and processes across different data sources. We specifically illustrate this approach for integrating LES and satellite data of aerosol-cloud interactions in subtropical stratocumulus cloud decks. We furthermore explore the extension of our framework to ground-based observations of Arctic mixed-phase clouds.

-----  
-

### References:

- Glassmeier, F., F. Hoffmann, J. S. Johnson, T. Yamaguchi, K. S. Carslaw and G. Feingold (2019): "An emulator approach to stratocumulus susceptibility", *Atmos. Chem. Phys.*, 19, 10191- 10203, doi: 10.5194/acp-19-10191-2019
- Hoffmann, F., F. Glassmeier, T. Yamaguchi and G. Feingold (2020): "Liquid water path steady states in stratocumulus: insights from process-level emulation and mixed-layer theory", *J. Atmos. Sci.*, 77, 2203-2215, doi: 10.1175/JAS-D-19-0241.1
- Glassmeier, F., F. Hoffmann, J.S. Johnson, T. Yamaguchi, K. S. Carslaw, and G. Feingold (2021): "Aerosol-cloud climate cooling overestimated by ship-track data", *Science* 371, 485-489, doi: 10.1126/science.abd3980