



EMS Annual Meeting Abstracts  
Vol. 20, EMS2023-145, 2023, updated on 29 May 2024  
<https://doi.org/10.5194/ems2023-145>  
EMS Annual Meeting 2023  
© Author(s) 2024. This work is distributed under  
the Creative Commons Attribution 4.0 License.



## Developing a method to estimate the ammonia loss from fertilized cropland and the part of soil emission recaptured by the canopy, applying field scale bidirectional models and flux measurements

László Horváth<sup>1,2</sup>, Csilla Gombi<sup>1</sup>, Helga Huszár<sup>1,2</sup>, Zoltán Nagy<sup>3</sup>, Krisztina Pintér<sup>4</sup>, Anna Szabó<sup>1,2</sup>, Tünde Takács<sup>5</sup>, Péter Torma<sup>2,6</sup>, Eszter Tóth<sup>5</sup>, Tamás Weidinger<sup>7</sup>, Gábor Szabó<sup>1</sup>, and Zoltán Bozóki<sup>1,2</sup>  
<sup>1</sup>University of Szeged, Institute of Physics, Department of Optics and Quantum Electronics, Szeged, Hungary  
(horvathl@titan.physx.u-szeged.hu)

<sup>2</sup>ELKH-SZTE Research Group for Photoacoustic Monitoring of Environmental Processes, H-6720 Szeged, Hungary

<sup>3</sup>Department of Plant Physiology and Plant Ecology, Institute of Agronomy, Hungarian University for Agriculture and Life Sciences, H-2100 Gödöllő, Hungary

<sup>4</sup>ELKH-MATE Agroecology Research Group, Hungarian University for Agriculture and Life Sciences, H-2100 Gödöllő, Hungary

<sup>5</sup>Department of Soil Physics and Water Management Institute for Soil Sciences and Agricultural Chemistry, Centre for Agricultural Research, H-1022 Budapest, Hungary

<sup>6</sup>National Laboratory for Water Science and Water Security, Budapest University of Technology and Economics, Faculty of Civil Engineering, Department of Hydraulic and Water Resources Engineering, H-1111 Budapest, Hungary

<sup>7</sup>Department of Meteorology, Institute of Geography and Earth Sciences Eötvös Loránd University, H-1117 Budapest, Hungary

In parameterization of the bidirectional ammonia exchange models over vegetated surfaces there are some crucial parameters: the stomatal, the soil, and the cuticular compensation point concentrations as the function of  $[\text{NH}_4^+]/[\text{H}^+]$  ratio in the apoplast, soil, and droplets on leaves, as well as the cuticular resistance. These factors determine the direction and the magnitude of the ammonia flux. Two-layer bidirectional exchange models are generally used to partition the measured flux into different parts. However, the parameterizations are mostly based on empirical relationships involving uncertainties and resulting in disagreements among the applied models in the estimation of the stomatal/soil/cuticular flux ratio. The main reasons for the deviations may be the following: i) Overestimation of the soil compensation-point when calculated from the bulk ammonium content of the soil because a part of ammonium content in the soil is bound in the solid phase. Hence Henry's law for the liquid phase cannot be applied to this fraction. ii) Neglecting the part of soil-derived ammonia recaptured by leaves. For this reason, soil emissions may be underestimated. iii) Lack of bioassay measurement. The models generally use empirical approximations to calculate the stomatal compensation point concentration e.g., by deriving it from the bulk ammonium content of the leaf tissue, which can be a source of bias. iv) Inaccurate or rough estimate of cuticular resistance, which is determined by the ratio of acidic air components and ammonia, besides the temperature and humidity. Models often consider a constant site-specific average for this parameter, even though the ratio of acidic substances to ammonia gas has diurnal and annual variations. Due to these uncertainties, the estimation of the

share of fluxes controlled by soil and vegetation is often uncertain. Furthermore, the uncertainty of the parameterization limits the model's applicability and reduces its robustness. As a conception, we aim to simultaneously measure the ammonia flux above the canopy and on the soil. Hence, as the difference between the two fluxes we can determine the amount of ammonia recaptured by the canopy. By means of the soil flux measurement, we can also estimate the bias of empirical soil flux parameterization from soil bulk ammonium content. We also plan performing bioassay measurements to exclude the potential error derived from empirical estimation. Also, we intend calculate the acid/base gas ratio using daily mean concentrations, considering the diurnal variations of humidity-dependent nitric acid/ammonia/ammonium nitrate equilibrium. We will use different conceptual models separately, based on previously developed simulation for all variations of day/night, wet/dry bare soil/vegetation cases in flux estimation at a fertilized crop, making it possible to partition the ratio of ammonia emitted by soil and recaptured either by stomata (built up in plant tissue) or by wet cuticula (net loss).