



## Evolution of surface changes and weather on Titan in the wake of the Fall 2010 equatorial storm

E.P. Turtle (1), J.E. Perry (2), A.G. Hayes (3), R.D. Lorenz (1), J.W. Barnes (4), A.S. McEwen (2), R.A. West (5), A.D. Del Genio (6), J.M. Barbara (6), J.I. Lunine (7), E.L. Schaller (2), T.L. Ray (5), R.M.C. Lopes (5), E.R. Stofan (8)  
(1) JHU/APL, Laurel, MD, USA (Elizabeth.Turtle@jhuapl.edu), (2) Univ. Arizona, Tucson, AZ, USA, (3) Caltech, Pasadena, CA, USA, (4) Univ. Idaho, Moscow, ID, USA, (5) JPL, Pasadena, CA, USA, (6) NASA GISS, New York, NY, USA, (7) INAF, Rome, Italy, (8) Proxemy Research, Rectortown, VA, USA.

Post-equinox changes in atmospheric circulation brought clouds and extensive methane rain to Titan's low latitudes, and large low-latitude clouds observed in Sept.-Oct. 2010 (Fig. 1) were quickly followed by changes (Fig. 2) attributed to extensive methane rainfall darkening the surface [1, 2]. Cassini Imaging Science Subsystem (ISS) observations in Oct. 2010 (Fig. 2) revealed differences in surface brightness along the southern boundary of Belet, one of Titan's largest dune fields. An area  $510,000 \pm 20,000 \text{ km}^2$ , extending  $\sim 2000 \text{ km}$  east-west and  $>130 \text{ km}$  across, darkened  $>10\%$ , adjacent areas remained unchanged.

It is likely that much of the darkening was caused by surface wetting [2], although runoff and ponding may have occurred in some areas, explaining the variation in the rates at which the surface has reverted to its original brightness as different areas drain (by overland flow or infiltration) or dry at different rates. A few small areas that appear to have brightened relative to their original appearance could be surfaces (potentially water ice [3]) cleaned by runoff or persistent low-altitude clouds or fog.

The frequency of and amount of precipitation from large storms at low latitudes during the equinoctial transition have important implications for Titan's methane cycle, atmospheric circulation, and rates of geologic modification. Although there is evidence that liquids have flowed on Titan's surface at the equator in the past -- fluvial channels are observed at all latitudes [4] -- to date liquids have only been observed on the surface at polar latitudes [5]. The Huygens Probe also detected methane moisture in the shallow subsurface [6-8] of the cobble-strewn flood plain at  $\sim 10^\circ\text{S}$  where it landed [9, 10]. Vast equatorial areas of long-lived longitudinal dunes [11] indicate that low latitudes are predominantly arid [12]; models predict low-latitude storms around equinox provide insufficient precipitation to accumulate there over the course of a year [13].

A series of images over the several months that have elapsed since the storm has revealed that in most of this area the changes have been short-lived: only a few patches persisted through April 2011 (Fig. 2). In an unsaturated permeable medium, vertical infiltration rates will be high ( $>20 \text{ mm/week}$  [14]), so persistence of surface liquids over a timescale of several months strongly suggests either a shallow impermeable layer or that the local methane table lies close to the surface. Evaporation rates of  $>1 \text{ mm/week}$  are predicted in equatorial regions [13] (rates of  $20 \text{ mm/week}$  have been documented at Titan's poles [15]), thus areas where darkening has persisted must be saturated ground at the methane table or have had ponded liquid of  $2.5\text{-}50 \text{ cm}$  depth.

We have continued to monitor Titan frequently (at least a few times per month), but to date no significant cloud activity has been observed since Fall 2010. This lack of clouds may indicate that the outbreak removed enough  $\text{CH}_4$  from the atmosphere and the lapse rate stabilized sufficiently that activity will not resume until the onset of convection at mid-northern latitudes later in the spring.




Figure 1: Extensive near-equatorial cloud, 27 Sept. 2010. Bright features are tropospheric methane clouds; grey and darker shades are surface features.




Figure 2: Sequence of ISS observations. Single arrow indicates isolated unchanged area between Belet and the darkened swath. Triple arrow highlights darkened swath on 29 Oct. 2010 and evolution thereof. Bright features on 27 Sept. and 29 Oct. are tropospheric methane clouds; grey and darker shades are surface features. North is up.

## Acknowledgements

We thank all who developed and operate the Cassini-Huygens mission. Research was supported by the Cassini-Huygens project.

## References

- [1] Turtle et al., Seasonal Changes in Titan's Meteorology, *GRL* 38, L03203, doi:10.1029/2010GL046266, 2011.
- [2] Turtle et al., Rapid and Extensive Surface Changes Near Titan's Equator: Evidence of April Showers, *Science* 331, p. 1414, 10.1126/science.1201063, 2011.
- [3] Clark et al., Detection and mapping of hydrocarbon deposits on Titan, *J. Geophys. Res.* 115, E10005, 2010.
- [4] Lorenz et al., Fluvial channels on Titan: Initial Cassini RADAR observations, *Planet. Space Sci.* 56, p. 1132, 2008.
- [5] Lopes et al., Distribution and interplay of geologic processes on Titan from Cassini radar data, *Icarus* 205, p. 540, 2010.
- [6] Karkoschka and Tomasko, Rain and dewdrops on titan based on in situ imaging, *Icarus* 199, p. 442, 2009.
- [7] Lorenz et al., Titan's damp ground: Constraints on Titan surface thermal properties from the temperature evolution of the Huygens GCMS inlet, *Meteorit. Planet. Sci.* 41, p. 1705, 2006.
- [8] Niemann et al., The abundances of constituents of Titan's atmosphere from the GCMS instrument on the Huygens probe, *Nature* 438, p. 779, 2005.
- [9] L. Soderblom et al., Topography and geomorphology of the Huygens landing site on Titan, *Planet. Space Sci.* 55, p. 2015, 2007.
- [10] M. G. Tomasko et al., Rain, winds and haze during the Huygens probe's descent to Titan's surface, *Nature* 438, p. 765, 2005.
- [11] Lorenz and Radebaugh, Global pattern of Titan's dunes: Radar survey from the Cassini prime mission, *Geophys. Res. Lett.* 36, L03202, 2009.
- [12] Lorenz et al., The Sand Seas of Titan: Cassini RADAR Observations of Longitudinal Dunes, *Science* 312, p. 724, 2006.
- [13] Mitchell, The drying of Titan's dunes: Titan's methane hydrology and its impact on atmospheric circulation, *Geophys. Res.* 113, E01805, 2008.
- [14] Hayes et al., Hydrocarbon lakes on Titan: Distribution and interaction with a porous regolith, *Geophys. Res. Lett.* 35, L09204, 2008.
- [15] Hayes et al., Transient surface liquid in Titan's polar regions from Cassini, *Icarus* 211, p. 655, 2011.