

# Condensed-Phase Photochemical Processes in Titan's Aerosols and Surface: The Role of Longer Wavelength Photochemistry

Murthy S. Gudipati (1), Ronen Jacovi (1), Antti Lignell (1), and Isabelle Couturier (2)  
(1) Science Division, Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109 USA, (gudipati@jpl.nasa.gov), (2) Université de Provence, 13397 Marseille cedex 20 France

## Abstract

We will discuss photochemical properties of Titan's organic molecules in the condensed phase as solid aerosols or surface material, from small linear polyynes (polyacetylenes and polycyanoacetylenes) such as  $C_2H_2$ ,  $C_4N_2$ ,  $HC_3N$ , etc. In particular we will focus on photochemistry caused by longer wavelength UV-VIS photons ( $>250$  nm) photons that make it through Titan's atmosphere to the haze region ( $\sim 100$  km) and on to the surface of Titan.

## 1. Introduction

Photochemistry of nitrogen and methane dominated Titan's atmosphere results in the formation of larger organics [1-5] that form the haze region in its lower atmosphere ( $<100$  km) [6, 7] and finally rain down on to Titan's surface [8, 9]. It is assumed that photochemistry is only confined to the upper atmosphere, where high-energy UV photons, electrons, Saturn's magnetospheric particles, and solar wind make their way through. At lower altitudes, due to the lack of high-energy radiation source (other than cosmic rays), it is expected that no further photochemistry could occur. Recent data from Cassini-Huygens [10] indicates that solar light at wavelengths  $>350$  nm makes its way through the haze region on to the surface (modulated by Titan's methane atmosphere). Thus, it is important to understand whether these UV-VIS photons can induce photochemistry in Titan's atmosphere, in particular in condensed-phase organics present in the atmosphere as well as on the surface.

## 2. Laboratory Experimental Studies

At the Titan's Organic Aerosol chemiSTry (TOAST) lab of JPL, we investigate spectroscopic and

photochemical properties of Titan's atmospheric molecules such as cyanoacetylenes. As these molecules can't be transported easily, we built dedicated laboratory to synthesize these molecules on the spot and use them for spectroscopic and photochemical studies. We carried out laser irradiation studies using 532 nm, 355 nm, and 266 nm (Nd-YAG laser 2<sup>nd</sup>, 3<sup>rd</sup>, and 4<sup>th</sup> harmonics). IR and UV spectra are monitored simultaneously to follow the reaction kinetics and reaction products.

Our focus is to determine whether at these wavelengths photochemical reactions involving Titan's organic molecules in condensed aerosol haze or on the surface can occur or not. In this talk we will discuss theoretical predictions and experimental findings on the longer wavelength photochemical behavior of Titan's organics.

## Acknowledgements

This research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration. NASA funding through NASA Astrobiology Institute Node "Titan Prebiotic Chemistry" for carrying out the research and JPL's DRDF and R&TD funding for the infrastructure and conduct preliminary studies.

## References

1. Kim, S.J., et al., Clouds, haze, and  $CH_4$ ,  $CH_3D$ ,  $HCN$ , and  $C_2H_2$  in the atmosphere of Titan probed via 3  $\mu m$  spectroscopy. *Icarus*, 2005. 173(2): p. 522-532.
2. Bezard, B., Composition and chemistry of Titan's stratosphere. *Philosophical Transactions of the Royal Society a-Mathematical Physical and Engineering Sciences*, 2009. 367(1889): p. 683-695.

3. Hirtzig, M., et al., A review of Titan's atmospheric phenomena. *Astronomy and Astrophysics Review*, 2009. 17(2): p. 105-147.
4. Gu, X., et al., Chemical dynamics of triacetylene formation and implications to the synthesis of polyynes in Titan's atmosphere. *Proceedings of the National Academy of Sciences of the United States of America*, 2009. 106(38): p. 16078-16083.
5. Zwier, T.S. and M. Allen, Metastable diacetylene reactions as routes to large hydrocarbons in Titan's atmosphere. *Icarus*, 1996. 123(2): p. 578-583.
6. Tomasko, M.G., et al., Rain, winds and haze during the Huygens probe's descent to Titan's surface. *Nature*, 2005. 438(7069): p. 765-778.
7. Penteado, P.F., et al., Latitudinal variations in Titan's methane and haze from Cassini VIMS observations. *Icarus*, 2010. 206(1): p. 352-365.
8. Clark, R.N., et al., Detection and mapping of hydrocarbon deposits on Titan. *Journal of Geophysical Research-Planets*, 2010. 115: p. 28.
9. Zarnecki, J.C., et al., A soft solid surface on Titan as revealed by the Huygens Surface Science Package. *Nature*, 2005. 438(7069): p. 792-795.
10. Lavvas, P.P., A. Coustenis, and I.M. Vardavas, Coupling photochemistry with haze formation in Titan's atmosphere, Part II: Results and validation with Cassini/Huygens data. *Planetary and Space Science*, 2008. 56(1): p. 67-99.

