

Trends in Cometary Dust Tails in NEOWISE

Emily Kramer (1), Amy Mainzer (1), James Bauer (2), Yanga Fernandez (3), Tommy Grav (4), Joseph Masiero (1), Sarah Sonnett (4), Edward Wright (5), and the NEOWISE Team.

(1) Jet Propulsion Laboratory, California Institute of Technology, California, USA, (2) University of Maryland, Maryland, USA, (3) University of Central Florida, Florida, USA, (4) Planetary Science Institute, Arizona, USA, (5) University of California Los Angeles, California, USA. (Emily.Kramer@jpl.nasa.gov)

Abstract

Comets are among the most primitive objects in our Solar System, having undergone little alteration since their formation in the protoplanetary disk. Yet the comets have still experienced some changes due to different heating processes such as impacts and insolation [1]. These processes may have caused significant structural changes to the comet, possibly depleting the comet of volatiles at the same time. As a comet approaches perihelion, the thermal energy from the Sun warms the comet, causing volatiles to be released, and intermixed dust particles to be carried along with the escaping gas. The dust and gas form the characteristic tails for which comets are so well-known. Since the comets themselves are relatively pristine, the dust particles that comprise their tails are thought to be similar to particles in the protoplanetary disk. By studying the dust tails of comets in different populations (long-period comets [LPCs] and short-period comets [SPCs]), we can investigate the effect of heating on the structure of comets.

NEOWISE is the planetary-funded mission that utilizes data from the Wide-field Infrared Survey Explorer (WISE) spacecraft to detect and characterize moving objects. The WISE mission surveyed the sky in four infrared wavelength bands (3.4, 4.6, 12 and 22-micron) between January 2010 and February 2011 [2, 3]. During the course of the prime mission, over 160 comets were serendipitously observed, including 22 newly discovered comets. About 89 of the comets observed by NEOWISE displayed a significant dust tail in the 12 and 22-micron (thermal emission) bands, showing a wide range of activity levels and dust morphology. A sample of six comets are shown in Figure 1. Since the observed objects are a mix of about 1/3 LPCs and 2/3 SPCs, differences in their activity can be used to better understand the thermal evolution that each of these populations has undergone.




Figure 1: Montage of six comets observed by NEOWISE during the cryogenic mission, showing the variety of activity levels present in the data. The images are in 22-micron, and each image is 10' across.

1. Methods

Each comet was serendipitously observed multiple times by NEOWISE, and the individual images were stacked to increase the signal-to-noise ratio. For the comets that displayed a significant dust tail, we have estimated the sizes and ages of the particles using dynamical models based on the Finsen-Probstein method [4, 5]. The Finsen-Probstein method assumes that the motion of cometary dust particles is controlled only by solar radiation pressure and solar gravity, which can be parameterized by the ratio $\beta = F_{rad}/F_{grav}$. β can be thought of as a proxy for particle size, with larger β corresponding to larger particles. β is incorporated into the equation of motion that can be integrated to track the motion of particles with different β values.

For a selection of 40 comets, we have then compared these models to the data using a novel tail-fitting method that allows the best-fit model to be chosen analytically rather than subjectively [6, 7]. For comets that were observed multiple times by WISE,

the dust tail particle properties were estimated separately, and then compared.

2. Results

We find that the dust tails of both LPCs and SPCs are primarily comprised of \sim mm to cm sized particles, which were the result of emission that occurred several months to several years prior to the observations. The LPCs nearly all have strong dust emission close to the comet's perihelion distance, and the SPCs mostly have strong dust emission close to perihelion, but some have strong dust emission well before perihelion. When comparing the two populations as a whole, we find that there is no statistical difference in the size of the particles between the two populations. Similar sized particles suggest similar internal structure and possibly a similar origin for the two populations. Further implications of these results will also be discussed.

3. Acknowledgements

This publication makes use of data products from the Wide-field Infrared Survey Explorer, which is a joint project of the University of California, Los Angeles and the Jet Propulsion Laboratory/California Institute of Technology, funded by the National Aeronautics and Space Administration. This publication also makes use of data products from NEOWISE, which is a project of the Jet Propulsion Laboratory/California Institute of Technology, funded by the Planetary Science Division of the National Aeronautics and Space Administration.

References

- [1] Prialnik, D. and Bar-Nun, A. (1987) *ApJ*, 313:893-905
- [2] Wright, E.L. et al. (2010) *AJ*, 140
- [3] Mainzer, A. et al. (2011) *ApJ*, 731:1
- [4] Finson, M. and Probstein, R. (1968) *ApJ*, 154
- [5] Lisse, C.M. et al. (1998) *ApJ*, 496
- [6] Kramer, E.A. (2014) PhD diss.
- [7] Kramer, E.A. et al. (2017) *ApJ*, 838