

Dynamics of slow ejecta from Comet 67P/Churyumov–Gerasimenko and the Places of their Depositions

L. Czechowski¹ and K.J. Kossacki².

¹University of Warsaw, Faculty of Physics, Institute of Geophysics, ul. Pasteura 5, 02-093 Poland
(lczech@op.pl).

²University of Warsaw, Faculty of Physics, Institute of Geophysics, ul. Pasteura 5, 02-093 Poland
(kjkossac@fuw.edu.pl).

Abstract

Comets have weak gravity field, so it is believed that probability of landslides is very low. However observations from space missions to comets 9P/Tempe 1 and 67P/Churyumov – Gerasimenko revealed existence of these mass motion.

In the present paper we consider comet 67P/Churyumov–Gerasimenko. Investigation of its nucleus indicated existence of deposits typical for landslides or/and deposition of slow ejecta.

There are some similarity of slow ejecta (i.e., with velocity lower than escape velocity) and landslides. Both are forms of gravity movement. After landing ejecta can be still moving like a ‘regular’ landslide. On the other hand, the motion of landslides can include free falling without contact with the ground ('ballistic' landslides).

The causes of landslides are usually related to instabilities of slopes. It is often possible to indicate a few causes of the landslide but usually only one factor is considered to be a trigger [1].

1. Topography of the surface

On the small bodies of spherical shape the area with large slope (in respect to the local gravity) is rather limited. The different situation could be observed on highly asymmetric bodies [2].

The gravitational field of 67P/Churyumov–Gerasimenko comet is very complicated. There are several regions of different slopes of the physical surface in respect to the gravity. The most of the surface ($\sim 74\%$) has the slope in the range $0 < \alpha < 40^\circ$.

The slope in the range $40^\circ < \alpha < 70^\circ$ is found on $\sim 17\%$ of the surface.

2. Origin of slow ejecta

A simple model of processes leading to the formation of slow ejecta is assumed. The phase transition heats a certain underground volume [3, 4]. It leads to vaporization of volatiles. Eventually a cavity is formed. If the pressure in the cavity exceeds some critical value then the crust could be crushed and its fragments will be ejected in space. Some parts of depression in Imhotep region (on the large lobe of the comet) could be a result of such processes.

3. Dynamics and deposition

In paper [5] slow ejecta from depression Hathemit (on the small lobe) are discussed. We found that ejecta with the velocity 0.3 m s^{-1} (or lower) land close to the starting point. Ejecta faster than 0.5 m s^{-1} have complex trajectories and could land far from the starting point. For the velocity in the range $0.5 – 0.6 \text{ m s}^{-1}$ the most of the ejecta from depression Hatmehit falls in a wide belt on one comet's hemisphere. These results are confirmed by observations: southern and northern hemisphere surfaces are clearly different [6].

In the present research we discuss slow ejecta from the large lobe. Fig. 1 presents 11 trajectories of test particles ejected with velocity 0.6 m s^{-1} from different places on this lobe. Most of them eventually have landed also on the same lobe.

We concentrated on ejecta from region of Imhotep. The large morphological changes was observed in this region. These changes could give rise to slow ejecta. The trajectories of test particles ejected with

the velocity 0.7 m s^{-1} from different parts of Imhotep are given in Fig. 2. Note that ejecta are deposited mainly in two different regions, one in the large lobe and another in the small lobe.




Fig. 1. Assumed mass distribution in the comet (green volume) and the trajectories of motion of the matter (red lines) ejected vertically (in respect to the physical surface) from 11 positions on the large lobe (right hand part on the figure) with the velocity 0.6 m s^{-1} . Note that the starting points of ejecta are on the upper surface of the lobe. Most of the landing points are on the upper surface on the large or small lobes. However, some trajectories reach also other sides of the comet.

Ejecta landing on the highly inclined surface could trigger another landslide. It depend on angles of landing and the properties of the material of the comet. In [5] we indicate that majority of the place of deposition are probably stable.

Determining the places of deposition of the material ejected from Imhotep or Hatmelib will allow to determine the composition of the comet's interior under these regions without the need for drilling. This would be particularly important if the CEASAR mission will be launched. It will return a sample from the nucleus.




Fig. 2. The trajectories of motion of the matter ejected from Imhotep (on the large lobe) with the velocity 0.7 m s^{-1} . Note that deposition of ejecta are concentrated mainly in two regions.

Acknowledgements

The research is partly supported by Polish National Science Centre (decision 2014/15/B/ST 10/02117).

References

- [1] Czechowski L., (2016) Earthquakes on comets and their consequences. LPSC 2016, 2781 pdf.
- [2] Czechowski L., (2017) Dynamics of landslides on comets of irregular shape. Geophysical Research Abstract. EGU 2017 April, 26, 2017
- [3] Kossacki K., Czechowski L., 2018. Comet 67P/Churyumov–Gerasimenko, possible origin of the depression Hatmehit. Icarus vol. 305, pp. 1-14, doi: 10.1016/j.icarus.2017.12.027
- [4] Kossacki, K.J., Szutowicz, S., 210, Crystallization of ice in Comet 17P/Holmes: Probably not responsible for the explosive 2007 megaburst. Icarus 207, 320- 340.
- [5] Czechowski L. and Kossacki K.J. 2019. Dynamics of material ejected from depression Hatmehit and landslides on comet 67P/Churyumov–Gerasimenko. Submitted.
- [6] Jui-Chi Lee, et al. 2016. Geomorphological mapping of comet 67P/Churyumov–Gerasimenko's Southern hemisphere. MNRAS 462, S573–S592 (2016) doi:10.1093/mnras/stx450