

# CLUPI, Close-UP Imager on the rover of the ExoMars mission 2020 to discover biofabrics on Mars - Science objectives and Science calibration

J.-L. Josset<sup>1</sup>, F. Westall<sup>2</sup>, B. A. Hofmann<sup>3</sup>, J. G. Spray<sup>4</sup>, C. Cockell<sup>5</sup>, S. Kempe<sup>6</sup>, A. D. Griffiths<sup>7</sup>, M.C. De Sanctis<sup>8</sup>, L. Colangeli<sup>9</sup>, D. Koschny<sup>9</sup>, T. Bontognali<sup>1</sup>, K. Föllmi<sup>10</sup>, E. Verrecchia<sup>11</sup>, L. Diamond<sup>12</sup>, M. Josset<sup>1</sup>, E. Javaux<sup>13</sup>, F. Esposito<sup>14</sup>, M. Gunn<sup>15</sup>, L. Fayon<sup>1</sup>, O. Korablev<sup>16</sup>, S. Erkman<sup>17</sup>, G. Paar<sup>18</sup>, S. Ulamec<sup>19</sup>, F. Foucher<sup>2</sup>, N. Kuhn<sup>20</sup>, J.-P. De Vera<sup>21</sup>, M. Tanevski<sup>1</sup>, P. Martin<sup>22</sup>; A. Toni<sup>9</sup>; J. Vago<sup>9</sup>

<sup>1</sup>Space Exploration Institute, Neuchâtel, Switzerland (jean-luc.josset@space-x.ch); <sup>2</sup>Centre de Biophysique Moléculaire, Orléans, France; <sup>3</sup>Natural History Museum, Bern, Switzerland; <sup>4</sup>Planetary and Space Science Centre, University of New Brunswick, Canada; <sup>5</sup>UK Center for Astrobiology, University of Edinburgh, Scotland; <sup>6</sup>Geosciences University of Technology Darmstadt, Germany; <sup>7</sup>University College London, United Kingdom; <sup>8</sup>Istituto di Astrofisica e Planetologia Spaziali, Roma, Italy; <sup>9</sup>ESA, The Netherlands; <sup>10</sup>Institute of Geology and Paleontology, University of Lausanne, Switzerland; <sup>11</sup>Institute of Earth Surface Dynamics, University of Lausanne, Switzerland; <sup>12</sup>Institute for Geological sciences, University of Bern, Switzerland; <sup>13</sup>Departement de Géologie, Unité PPM, University of Liège, Belgium; <sup>14</sup>Osservatorio Astronomico di Capodimonte, Napoli, Italy; <sup>15</sup>Aberystwyth University, United Kingdom; <sup>16</sup>IKI, Space Research Institute, Moscow, Russia; <sup>17</sup>Faculty of Geosciences and Environment, University of Lausanne, Switzerland; <sup>18</sup>Joanneum Research, Graz, Austria; <sup>19</sup>DLR, Space Operations, Cologne, Germany; <sup>20</sup>Physical Geography and Environmental Change, University of Basel, Switzerland; <sup>21</sup>DLR, Institute of Planetary Research, Berlin, Germany; <sup>22</sup>LPCEE, Orléans, France.

## Abstract

CLUPI, the high-performance colour close-up imager, on board the 2020 ExoMars Rover plays an important role in attaining the mission objectives: it is the equivalent of the hand lens that no geologist is without when undertaking field work. CLUPI is a powerful, highly integrated miniaturized (<900g) low-power robust imaging system, able to sustain very low temperatures (−120°C). CLUPI has a focus mechanism allowing a working distance from 11.5cm to infinite providing outstanding pictures with a color detector of 2652x1768x3. At 11.5cm, the spatial resolution is 8 micrometer/pixel in color. The concept benefits from well-proven heritage: Proba, Rosetta, MarsExpress and Smart-1 missions...

In a typical field scenario, the geologist will use his/her eyes to make an overview of an area and the outcrops within it to determine sites of particular interest for more detailed study. In the ExoMars scenario, after having made a preliminary general evaluation, the geologist will approach a particular outcrop for closer observation of structures at the decimetre to subdecimeter scale before finally getting very close up to the surface with a hand lens (CLUPI), and/or taking a hand specimen, for detailed

observation of textures and minerals. Using structural, textural and preliminary compositional analysis, the geologist identifies the materials and makes a decision as to whether they are of sufficient interest to be subsampled for laboratory analysis (using the ExoMars drill and laboratory instruments).

Given the time and energy expense necessary for drilling and analysing samples in the rover laboratory, preliminary screening of the materials to choose those most likely to be of interest is essential. ExoMars will be choosing the samples exactly as a field geologist does – by observation (backed up by years and years of field experience in rock interpretation in the field). Because the main science objective of ExoMars concerns the search for life, whose traces on Mars are likely to be cryptic, close up observation of the rocks and granular regolith will be critical to the decision as to whether to drill and sample the nearby underlying materials. Thus, CLUPI is the essential final step in the choice of drill site. But not only are CLUPI's observations of the rock outcrops important, but they also serve other purposes. CLUPI, could observe the placement of the drill head. It will also be able to observe the fines that come out of the drill hole, including any colour stratification linked to lithological changes with depth. Finally, CLUPI will

provide detailed observation of the surface of the core drilled materials when they are in the sample drawer at a spatial resolution of about 17 micrometer/pixel in color.

After a brief review of the science objectives of CLUPI together with the observation modes of the instrument, the science calibration performed on the flight model will be described, showing its importance for CLUPI capabilities to provide important information significantly contributing to the understanding of the geological environment and could identify outstanding potential biofabrics of past life on Mars.