

***Ab initio* reaction rate coefficients relevant to Titan conditions and cool exoplanet atmospheres**

Shiblee R. Barua^{1,2}, Paul N. Romani¹

¹NASA Goddard Space Flight Center, Greenbelt, Maryland USA (shiblee.barua@nasa.gov)

²Universities Space Research Association, Columbia, Maryland, United States

Existing photochemical models for the atmospheres of cool planetary bodies suffer from large errors when predicting the mole fraction profiles of various compounds. Global sensitivity analyses show that a major source of these model errors can be attributed to inaccurate rate coefficients for “key” radical–radical and radical–neutral reactions in low-temperature conditions. Accurate rate coefficients for key reactions thus improve the accuracy of photochemical models. Unfortunately, accurate experimental rate coefficients for such low- T reactions are extremely difficult, if not impossible, to measure since the lab data are affected by uncertainties in determining the absolute concentrations of the radical species. Currently, the most common theoretical approach involves uncertainty extrapolation technique in which uncertainties in room-temperature rate coefficients are extrapolated to low-temperature conditions, resulting in large errors in the theoretical low- T rate coefficient data. To solve this existing problem, we are employing the two-transition-state (2TS) model developed by Klippenstein and coworkers to calculate high-level *ab initio* rate-coefficients for key low- T (*i.e.*, 10–200 K) reactions relevant to Titan conditions and other possibly similar cool exoplanetary bodies. In particular, we are investigating key reactions that have not yet been studied in the lab, and for which accurate rate coefficients are still unknown. We will make our calculated *ab initio* rate coefficients available to the astrochemistry community via well-established free online kinetic databases (*e.g.* PDS, KIDA). Photochemical modelers will use these low- T rate coefficients to make accurate mole fraction predictions for cool planetary atmospheres, thus improving our understanding of the diverse chemistry of these bodies.

Table 1: Selected key reactions currently being investigated that are responsible for uncertainties in the mole fractions of major nitrogen species (left-most column) in low- T Titan conditions. Each reaction shown here has an absolute Rank Correlation Coefficient (RCC) value, a quantitative indicator of the sensitivity of model mole fractions on rate coefficients, greater than the chosen threshold of 0.2 at 100 km or above (right-most column).

Affected N Species	Key Reactions	 RCC > 0.2
NH ₃	$\text{N}({}^4\text{S}) + \text{NH}_2 \rightarrow \text{N}_2 + \text{H} + \text{H}$	(100, 900) km
HNC	$\text{N}({}^4\text{S}) + {}^3\text{CH}_2 \rightarrow \text{HNC} + \text{H}$	100 km
CH ₂ NH	$\text{N}({}^2\text{D}) + \text{CH}_4 \rightarrow \text{NH} + \text{CH}_3$	900 km
CH ₃ C ₃ N	${}^1\text{CH}_2 + \text{CH}_4 \rightarrow \text{CH}_3 + \text{CH}_3$	900 km
CH ₃ C ₃ N	$\text{C}_2\text{N} + \text{H} \rightarrow \text{HCN} + \text{C}$	900 km
C ₂ H ₃ CN	$\text{CHCN} + \text{H} \rightarrow \text{C}_2\text{N} + \text{H}_2$	900 km




Figure 1. Schematic one-dimensional picture of the potential energy as a function of the reaction coordinate for a neutral-neutral reaction leading to bimolecular products. In its approach, the 2TS model incorporates both the 1st (outer) transition state for barrierless association reaction and the 2nd (inner) transition state for reaction path with a barrier.