

The ultra-cold night sides of the hot and super-hot Jupiters WASP-43b and WASP-18b, arising with deep wind flow

Ludmila Carone (1), Robin Baeyens (2), Paul Mollière (3), Patrick Barth (1), Allona Vazan (4,5), Leen Decin (2), Paula Sarkis (6), Olivia Venot (7) and Thomas Henning (1)
(1) MPIA Heidelberg, Germany (carone@mpia.de), (2) KU Leuven, Belgium, (3) Leiden Observatory, The Netherlands, (4) The Hebrew University, Israel, (5) University of Zürich, Switzerland, (6) University of Bern, Switzerland, (7) Laboratoire Interuniversitaire des Systemes Atmospheriques, Paris, France

Abstract

We have shown that the hot Jupiter WASP-43b may develop strong westward winds at the equatorial day side, that is, anti-rotation - if deep wind flow ($p > 10$ bar) is taking into account [1]. Because equatorial anti-rotation prevents efficient horizontal heat transport, we also predict that the night side of WASP-43b will emit very little thermal flux (≤ 1000 ppm) for wavelengths 5 - 12 micron.

Here, we compare our 3D climate results for the hot Jupiter WASP-43b ($T_{\text{eff}}=1400$ K) with new results for the super-hot Jupiter WASP-18b ($T_{\text{eff}}=2400$ K) that was recently shown to also have inefficient horizontal heat distribution [2]. In the work of [2], the observational signals were explained by imposing magnetic drag throughout the atmosphere, which suppresses zonally banded jets and leads to direct or radial day-to-night-side flow. We will show an alternative climate scenario for WASP-18b with (partially) banded wind flow in our 3D GCM with deep wind flow.

We predict that, JWST/MIRI observations of the thermal emission at the night side of WASP-43b and WASP-18b will unambiguously distinguish between a climate scenario with wind flow at depth and ultra-cool night sides and other scenarios (fully superrotating jet with clouds [3] and drag-induced climate without jets [2]).

References

- [1] Carone, L. et al.: Equatorial anti-rotating day side wind flow in WASP-43b elicited by deep wind jets?, eprint arXiv:1904.13334, 2019.
- [2] Arcangeli, J. et al.: Climate of an Ultra hot Jupiter: Spectroscopic phase curve of WASP-18b with HST/WFC3, eprint arXiv:1904.02069, 2019.

- [3] Parmentier, V. et al.: Transitions in the Cloud Composition of Hot Jupiters, A & A, Vol. 828, Issue 1, article id. 22, 2016.