

FTIR and X-ray micro-tomography on Itokawa particles and carbonaceous chondrites

R. Brunetto¹, Z. Dionnet^{1,2}, A. Aléon-Toppani¹, D. Baklouti¹, F. Borondics³, Z. Djouadi¹, A. King³, T. Nakamura⁴, S. Rubino^{1,3}, C. Sandt³, D. Troadec⁵ and A. Tsuchiyama⁶

¹Institut d'Astrophysique Spatiale, CNRS, Université Paris-Saclay, Orsay, France (rosario.brunetto@ias.u-psud.fr), ²DIST-Università Parthenope, Napoli, Italy, ³SOLEIL synchrotron, Saint-Aubin, France, ⁴Division of Earth and Planetary Materials Science, Graduate School of Science, Tohoku University, Japan, ⁵Institut d'Electronique, de Microélectronique et de Nanotechnologie, Lille, France, ⁶Research Organization of Science and Technology, Ritsumeikan University/Guangzhou Institute of Geochemistry

Abstract

We report a combined IR and X micro-tomography analysis of five particles from asteroid Itokawa and a few selected fragments of several carbonaceous chondrites. These techniques are an efficient first-step in a multi-technique analytical sequence to be applied on small Solar System bodies samples, collected by space missions such as Hayabusa2 and OSIRIS-REx.

1. Introduction

Laboratory analyses of meteorites and interplanetary dust particles originating from asteroids and comets give us the opportunity to study directly the components that formed in the protoplanetary disk, but they also have intrinsic limits and biases, some of which are overcome by sample return space missions [1]. In the near future a new generation of sample return missions (Hayabusa2, OSIRIS-REx, MMX, etc.) will collect samples from small Solar System bodies. To maximize the scientific outcome of laboratory studies and minimize the loss of precious extraterrestrial samples, an analytical sequence from less destructive to more destructive techniques needs to be established. Among the possible techniques, infrared (IR) spectroscopy is important for being totally non-destructive and comparable to remote sensing observations of small bodies [2]. Thanks to IR imaging microspectroscopy, we are able to study the spatial distribution of molecular bonds associated to minerals, water and organics [3].

2. Method

In this work, we present the results of a Fourier-Transform infrared (FTIR) micro-tomography study of five particles from asteroid Itokawa collected

during the Hayabusa mission (JAXA) and of several fragments of selected carbonaceous chondrites (CM Paris and Cold Bokkeveld, C2-ung Tagish Lake, CK NWA 5515). All samples size 20-50 μm and are mounted on needles. The FTIR measurements are performed at the SMIS beamline of synchrotron SOLEIL (France) with a spatial resolution that is diffraction-limited and a voxel size of about 0.66 μm . Complementary X-ray micro-tomography analyses are performed at the PSICHE beamline of SOLEIL and at beamline BL47XU of the SPring-8 synchrotron (Japan) [4] with voxel sizes of about 0.127 μm and 0.084 μm respectively.

3. Results

Supported by X-ray micro-tomography analyses providing a shape model and 3D structures of the samples, the FTIR analyses allows the detection of mineral phases, water and organics, and their spatial co-localization in three dimensions. We show that a combined IR and X-ray micro-tomography analytical approach is able to provide a non-destructive 3D physical and chemical characterization of individual extraterrestrial particles. The two techniques together surmount the intrinsic limitations of each one, giving access to the identification and spatial distribution of individual components inside the analyzed particles, as well as their 3D structure and porosity. In addition, we show that it is possible to recover the samples after the analyses.

4. Summary and Conclusions

We propose these techniques as an efficient first-step in a multi-technique analytical sequence on samples collected by space missions. In particular, we consider FTIR 3D micro-tomography an interesting starting point, as it will be able to provide a first quick look on the composition, abundance and 3D

distribution of carbonaceous materials and water at the scale of few microns within individual grains sizing up to several tens of microns. Once the organics are revealed by IR measurements, thin sliced sections of the samples can be prepared to be analyzed by more destructive techniques, in order to retrieve the structure and elemental/isotopic composition of the carbonaceous component and its mineral host, down to nm scale. This top-down sequence applied to Ryugu samples collected by the Hayabusa2 mission (JAXA), may build a bridge between the observations at macroscopic scale [5] and the chemical and physical processes operating at the nano-scale.

Acknowledgements

The FTIR microspectroscopy activities are supported by grants from Région Ile-de-France (DIM-ACAV) and SOLEIL. This work has been funded by the Centre National d'Etudes Spatiales (CNES-France, Hayabusa2 mission) and by the ANR project CLASSY (Grant ANR-17-CE31-0004-02) of the French Agence Nationale de la Recherche. We are grateful to the JAXA Curator for allocating the Hayabusa particles, and to B. Zanda, K. Nakamura-Messenger and C. Lantz for providing us with the meteorite samples. We thank T. Yada and L. Bonal for useful discussion, and the ANATOMIX team (SOLEIL) for their help with Avizo.

References

- [1] Brunetto R and Lantz C. (2019) *Nature Astronomy* 3:290–292.
- [2] Brunetto R. et al. (2011) *Icarus* 212:896–910.
- [3] Dionnet Z. et al. (2018) *Meteoritics and Planetary Science* 53:2608–2623.
- [4] Tsuchiyama A. et al. (2013) *Geochimica and Cosmochimica Acta* 116:5–16.
- [5] Kitazato K. et al. (2019) *Science* DOI:10.1126/science.aav7432