



Impact as a geological process in the context of solar system and planetary evolution

G. Schmidt (1)
(1) Benswood Studios, Germany (schmidtgerhard@aol.com)

Abstract

The nature of planetary surfaces, e.g. Moon, Mars, Mercury, and Earth, attest to the importance of impact as a geological process in the context of solar system and planetary evolution.

1. Introduction

Understanding the origin of solid planets (core, mantle and crust) it is important to know relative abundances of refractory highly siderophile elements (Os, Ir, Ru, Rh) in silicate mantle and crust of planets. Osmium and Ir are abundant in most meteorites but depleted in crustal rocks (low target/meteorite ratios) and thus the most reliable elements for projectile identification. Chromium isotopic studies (excepting carbonaceous chondrites [1]), Os isotopic studies and estimates of Au/(Os, Ir, Ru, Rh) ratios are not helpful in identifying the meteoritic types of impactors on Earth.

1.1 Target rocks

However, target/meteorite ratios are high if target rocks consist of mantle rocks. In such cases Os, Ir, Ru, and Rh element contents are enriched in impactites due to relatively high abundances (ng/g level) in target rocks to make the identification of projectile types difficult (e.g., Gardnos impact structure, Norway [2]). Ruthenium/Ir is the most reliable key ratio that rules out Earth primitive upper mantle (PUM) derived refractory highly siderophile element components in impactites. Ruthenium/Ir of the Earth mantle is 2.0 ± 0.1 and thus significantly above chondritic ratios varying from 1.4 to 1.6 [3,4].

2. Earth mantle

On Earth Rh/Ir, Ru/Ir, Pd/Ir, and Pt/Os derived from PUM match the ratios of group IV irons with fractionated trace element patterns. The question raise if HSE in mantle rocks are added to the accreting Earth by a late bombardment of pre-

differentiated objects or the cores of these objects (magmatic iron meteorites as remnants of the first planetesimals) or some unsampled inner solar system materials from the Mercury-Venus formation region, not sampled through meteorite collections [5].

3. Earth crust

The PGE and Ni systematics of the upper continental crust (UCC) closely resembles group IIIAB iron meteorites with highly fractionated refractory trace element patterns, pallasites, and the evolved suite of Martian meteorites (representing Martian crust).

4. Projectiles of impact craters

To date, about 20 iron meteorites and about 20 chondrites have been identified as projectiles of 176 known small and large impact craters on Earth. The projectiles for 136 impact craters are still unknown.

5. Summary and Conclusions

Melt rocks from many impact craters on Earth (e.g., Rochechouart [6], Säcksjärvi [7], Dellen, Mien, Boltysh) are depleted in Os relatively to Ir and Ru. Subchondritic Os/Ir (about 3 times lower than Os/Ir in the metal phase of non-magmatic irons), fractionated Ru/Ir and Rh/Os ratios are strong arguments to conclude that about 50% of the currently identified iron projectiles from terrestrial impact craters are related to magmatic irons.

References

- [1] Koeberl, C., Shukolyukov, A., and Lugmair, G. W.: Chromium isotopic studies of terrestrial impact craters: Identification of meteoritic components at Bosumtwi, Clearwater East, Lappajärvi, and Rochechouart, *Earth and Planetary Science Letters* 256, 534–546, 2007.
- [2] Goderis, S., Kalleson, E., Tagle, R., Dypvik, H., Schmitt, R.-T., Erzinger, J., and Claeys, P.: A non-magmatic iron projectile for the Gardnos impact event, *Chemical Geology* 258, 145-156, 2009
- [3] Schmidt, G.: Are high-temperature fractionations in the solar nebula preserved in highly siderophile element systematics of the Earth's mantle? *Meteorit. Planet. Sci.* 39, 1995–2007, 2004.
- [4] Becker, H., Horan, M.F., Walker, R.J., Gao, S., Lorand, J.-P., and Rudnick, R.L.: Highly siderophile element composition of the Earth's primitive upper mantle: Constraints from new data on peridotite massifs and xenoliths, *Geochimica et Cosmochimica Acta* 70, 4528–4550, 2006.
- [5] Schmidt, G.: Planet Formation in the Inner Solar System: Origin and Nature of Impacting Projectiles, *Workshop on Planet Formation and Evolution: The Solar System and Extrasolar Planets*, 2-6 March 2009, Tübingen, Germany, 2009.
- [6] Janssens, M.-J., Hertogen, J., Takahashi, H., Anders, E., and Lambert, P., Rochechouart meteorite crater: Identification of projectile, *Journal of Geophysical Research* 82, 750-758, 1977.
- [7] Schmidt, G., Palme, H., and Kratz, K.-L., Highly siderophile elements (Re, Os, Ir, Ru, Rh, Pd, Au) in impact melts from three European craters (Sääksjärvi, Mien and Dellen): clues to the nature of the impacting bodies, *Geochimica et Cosmochimica Acta* 61, 2977–2987, 1997.