



Incorporation of volatiles into clathrates on Pluto and Triton

O. Mousis (1), A. Moudens (2), J. I. Lunine (3), S. Picaud (1), C. Thomas (1) and H. A. Weaver (4)

(1) Institut UTINAM, CNRS-UMR 6213, Observatoire de Besançon, BP 1615, 25010 Besançon Cedex, France (olivier.mousis@obs-besancon.fr / Fax: +33 381 666 944), (2) Institut de Physique de Rennes, Université de Rennes 1, 35042 Rennes Cedex, France, (3) Dipartimento di Fisica, Università degli Studi di Roma "Tor Vergata", Rome, Italy, (4) Space Department, Johns Hopkins University Applied Physics Laboratory, 11100 Johns Hopkins Road, Laurel, MD 20723-6099, USA

Abstract

We use a statistical thermodynamic approach to determine the relative abundances of guests in clathrate hydrates which may form from a multiple compound gas whose composition is similar to those of the atmospheres of Pluto and Triton. We consider different values of the methane, carbon monoxide, nitrogen, argon, krypton and xenon abundances that may be representative of the early atmospheres of Pluto and Triton. We then investigate the trapping efficiencies of various compounds sequestered in clathrate hydrates assumed to be formed at the atmosphere-surface interfaces of these bodies as a function of seasonal changes and orbital evolution. We quantify the amount of the various gases trapped as a function of the amount of clathrate formed in the crust, and propose observational tests that will allow future spacecraft missions to assess the plausible presence of clathrate hydrates in these environments.