

# Interpretation of the D/H ratio in comets: implications for outer solar system formation

**O. Mousis** (1), J.-M. Petit (1) and J. J. Kavelaars (2)

(1) Institut UTINAM, CNRS / Université de Franche Comté, Besançon, France, (2) Herzberg Institute of Astrophysics, National Research Council of Canada, Victoria, Canada (olivier.mousis@obs-besancon.fr / Fax: +33 381 666 944)

## 1. Introduction

A comet's origins in the primitive nebula can be probed by examining the degree to which fossil deuterium is enriched compared to the protosolar abundance. Calculations of the temporal and radial evolution of the deuterium enrichment in the solar nebula can reproduce existing D/H measures for comets [1, 2, 3]. These calculations show that the deuterium enrichment in water ice strongly depends on the distance from the Sun at which the ice was formed. Comparing the D/H value measured in comets with those predicted by such models allows retrieval of their formation location.

The measurement of the D/H ratio at Enceladus by the Ion and Neutral Mass Spectrometer aboard the Cassini spacecraft [4] provides a new, and tighter, constraint on the deuterium enrichment profile in the outer solar nebula prompting us to reconsider models presented in previous works [5].

## 2. Reservoirs and Source Regions of Comets

The “cometary reservoir” is the region of semi-stable phase space from which comets are currently being delivered, while the “source regions” are those parts of the primitive nebula in which the comets formed and were then delivered to the reservoirs. Ecliptic and isotropic comets are being delivered from at least two distinct reservoirs and, as such, are likely from different source regions.

The reservoir of the isotropic comets is, generically, the Oort cloud (see [6] for a good review). Some fraction of the isotropic comets with  $a < 20,000$  AU may arrive from the “innermost” component of this distribution [7], the remainder coming from the outer Oort cloud. Modeling of delivery into the Oort cloud reservoir generally finds this process to be controlled by Uranus-Neptune scattering.

If Uranus and Neptune originated at (roughly) 12 and 15 AU then material currently being delivered from the Oort cloud reservoir should have originated from a source much closer to the Sun than in cases where Uranus and Neptune formed at or near their current locations ( $\sim 20$  & 30 AU). A tracer of the chemical evolution of the primordial solar system that is sensitive to variations in the physical conditions between 10 and 30 AU, an example of which is described in the next section, provides a discriminator between these formation scenarios.

## 3. D/H in the Solar Nebula

Figure 1 describes the evolution of  $f$ , defined as the ratio of D/H in  $\text{H}_2\text{O}$  to that in molecular  $\text{H}_2$ , as a function of distance from the Sun in the case of a typical solar nebula. As in previous work, we assume that  $f$  is constant at  $t = 0$  irrespective of the heliocentric distance and corresponds to the value measured in the highly enriched component found in LL3 meteorites ( $\text{D/H} = (73 \pm 12) \times 10^{-5}$  [8]) compared to the protosolar value ( $(2.1 \pm 0.4) \times 10^{-5}$  [9]). The highly enriched component in LL3 meteorites is presumed to originate from ISM grains that were not reprocessed when entering the nebula [2].

For the adopted set of parameters, the deuterium enrichment profile simultaneously matches the nominal D/H value measured in  $\text{H}_2\text{O}$  in the moderately enriched component of LL3 meteorites at 3 AU and at the current heliocentric distance of Saturn matches the D/H enrichment of Enceladus. We were unable, in this investigation, to find models matching both the moderately enriched component of the LL3 meteorites at 3 AU and the value at Enceladus at 10 AU that did not also require the value of  $f$  to increase to much larger values in the region beyond 15 AU. Thus, the result that  $f$  in the 20-30 AU zone should have exceeded  $\sim 25$  is a generic outcome of the temperature evolution of the disk, when constrained by the D/H measured at Enceladus, and not particularly dependent on

the model of that evolution.




Figure 1: Enrichment factor  $f$  as a function of the heliocentric distance. The dashed curves correspond to the evolution of  $f$  in the gas phase prior to condensation terminated by dots at the heliocentric distance where  $H_2O$  condenses at the given epoch. The solid curve represents the value of  $f$  acquired by ice as a function of its formation distance in the nebula.

## 4. Conclusions

1P/Halley, 8P/Tuttle, C/1995 O1 (Hale-Bopp), C/1996 B2 (Hyakutake), and C/2001 Q4 (NEAT) all have D/H values that are consistent with or slightly larger than that of Enceladus. These comets are all members of the nearly-isotropic class and are, thus, drawn from a reservoir in some part of the Oort cloud. Based on dynamical arguments, the Oort cloud itself was fed by material from the Uranus/Neptune region. Our modeling of the dependence of  $f$  (pinned by the measured deuterium enrichment of Enceladus) on formation location (see Figure 1) precludes these comets from having formed beyond  $<15$  AU from the Sun. This implies that Uranus and Neptune were originally closer to the current location of Saturn than observed today, a configuration close to that preferred in the Nice model.

103P/Hartley 2 being D-poor ( $D/H \sim \text{SMOW}$ ) compared to these bodies questions our model. An unlikely solution would be to assume that ecliptic comets formed closer to the Sun than nearly isotropic comets. Models with a more realistic presentation of the origin and evolution of the D/H ratio in  $H_2O$  could be also taken into account. Such models could include initial heating of the exterior of the disk by infalling envelope or local heating of the disk by vortices. An alternative possibility would be that the D/H measurement in the

cometary comae does not represent the bulk composition of the nucleus, as it is suggested by recent experiments [10].

## Acknowledgements

J.J. Kavelaars acknowledges support provided by Embassy France. O. Mousis and J.-M. Petit acknowledge support provided by the Centre National d'Etudes Spatiales.

## References

- [1] Drouart, A., Dubrulle, B., Gautier, D. & Robert, F.: Structure and Transport in the Solar Nebula from Constraints on Deuterium Enrichment and Giant Planets Formation, *Icarus*, 140, 129, 1999.
- [2] Mousis, O., Gautier, D., Bockelée-Morvan, D., Robert, F. *et al.*: Constraints on the Formation of Comets from D/H Ratios Measured in  $H_2O$  and HCN, *Icarus*, 148, 513, 2000.
- [3] Horner, J., Mousis, O. & Hersant, F.: Constraints on the Formation Regions of Comets from their D:H Ratios, *Earth Moon and Planets*, 100, 43, 2007.
- [4] Waite, J.H.Jr., Lewis, W.S., Magee, B.A., Lunine, J.I. *et al.*: Liquid water on Enceladus from observations of ammonia and  $^{40}\text{Ar}$  in the plume, *Nature*, 460, 487, 2009.
- [5] Kavelaars, J.J., Mousis, O., Petit, J.-M. & Weaver, H.A.: On the Formation Location of Uranus and Neptune as constrained by Dynamical and Chemical Models of Comets, *Astrophys. J. Letters*, 734, L30, 2011.
- [6] Dones, L., Weissman, P.R., Levison, H.F. & Duncan, M.J.: *Comets II*, University of Arizona Press, Tucson, 2004.
- [7] Kaib, N.A. & Quinn, T.: Reassessing the Source of Long-Period Comets, *Science*, 325, 1234, 2009.
- [8] Deloule, E., Robert, F. & Doukhan, J.C.: Interstellar hydroxyl in meteoritic chondrules: implications for the origin of water in the inner solar system, *Geochim. Cosmochim. Acta*, 62, 3367, 1998.
- [9] Geiss, J. & Gloeckler, G.: Abundances of Deuterium and Helium-3 in the Protosolar Cloud, *Space Science Reviews*, 84, 239, 1998.
- [10] Brown, R. H., Lauretta, D. S., Schmidt, B., Moores, J.: Experimental and theoretical simulations of ice sublimation with implications for the chemical, isotopic, and physical evolution of icy objects, *Planetary and Space Science* 60, 166, 2012.