

## Cold Ion Escape from the Martian Ionosphere

**M. Fränz** (1), E. Dubinin (1), Y. Wei (1), J. Woch (1), D. Morgan (2), S. Barabash (3), R. Lundin (3) and A. Fedorov (4)

(1) MPI for Solar System Research, Katlenburg-Lindau, Germany (fraenz@mps.mpg.de / Fax: +49-5556-979-240),

(2) Physics Department, University of Iowa, USA, (3) Institute for Space Physics, Kiruna, Sweden, (4) Institut de Recherche en Astrophysique et Planetologie, Toulouse, France (fraenz@mps.mpg.de)

It has always been challenging to observe the flux of ions with energies of less than 10eV escaping from the planetary ionospheres. We here report on new measurements of the ionospheric ion flows at Mars by the ASPERA-3 experiment on board Mars Express. We first use support from the MARSIS radar experiment for some orbits with fortunate observation geometry. Here we have observed a transterminator flow of  $O^+$  and  $O_2^+$  ions with a super-sonic velocity of around 5km/s and fluxes of  $0.8 \cdot 10^9/cm^2s$ . If we assume a symmetric flux around the terminator this corresponds to an ion flow of  $3.1 \pm 0.5 \times 10^{25}/s$  half of which is expected to escape from Mars (Fraenz et al, 2010). This escape flux is significantly higher than previously observed on the tailside of Mars, we discuss possible reasons for the difference. Since 2008 the MARSIS radar does nightside local plasma density measurement which often coincide with ASPERA-3 measurements. In a new analysis of the combined nightside datasets (Fig. 1) we show that the main escape channel is along the shadow boundary on the tailside of Mars. At a distance of about  $0.5 R_M$  the flux settles at a constant value (Fig. 2) which indicates that about half of the transterminator ionospheric flow escapes from the planet.

Possible mechanism to generate this flux can be the ionospheric pressure gradient between dayside and nightside or momentum transfer from the solar wind via the induced magnetic field since the flow velocity is in the Alfvénic regime.




Figure 1: Median oxygen ion flux reconstructed by combining ion velocity observations of the Mars Express ASPERA-3 IMA sensor and local plasma density observations by the MARSIS radar. Each bin value is the median from observations on about 3000 orbits between May 2007 and July 2011. Horizontal axis is MSO X-axis (Sun towards the left), vertical axis is vertical distance from MSO X-axis.




Figure 2: Ring median flux of cylindrical ring regions of all bins shown in previous figure. The different colors show median fluxes for different regions in R-cylindrical (distance from MSO-X-axis) as a function of tailward distance from the terminator (or surface for  $R_{Cyl} < 1$ ).