

# Anion Chemistry On Titan: Probing the Destruction Mechanisms of Nitrile Anions by Interaction with Photons

J. Zabka (1), M. Polášek (1), M. Bradyová (1), Z. Flenerová (1), M. Obluková (1), D. Ascenzi (2), V. Vuitton (3), A. Giuliani (4), L. Nahon (4), A. R. Milosavljevic (5), C. Romanzin (6), C. Alcaraz (6)

(1) J. Heyrovsky Institute of Physical Chemistry of the ASCR, v. v. i. 182 23 Prague 8, Czech Republic, (2) Dept of Physics, Uni. Trento, Via Sommarive 14, 38050 Povo (TN), Italy, (3) Institut de Planétologie et d'Astrophysique de Grenoble, UMR 5274, BP 53, 38041 Grenoble, France, (4) Synchrotron SOLEIL, L'Orme des Merisiers, Saint Aubin BP 48, 91192 Gif sur Yvette Cedex, France, (5) Lab. for Atomic Collision Processes, Inst. of Physics, Uni. Belgrade, 11080 Zemun, Belgrade, (6) Lab. for Atomic Collision Processes, Inst. of Physics, Uni. Belgrade, 11080 Zemun, Belgrade, Serbia

## Abstract

The aim of this work is to study the interaction with VUV photons of mass-selected negative ions relevant for the understanding of Titan atmosphere. Characterization of their formation [1] and destruction rate is of fundamental importance for modeling Titan ionosphere chemistry and understanding the observations of heavy anions by the CAPS/ELS spectrometer on board of the CASSINI spacecraft. The objective here is to measure their transformation into smaller anions through photodissociation and their destruction through photodetachment. The parent anions  $CN^-$  are produced from  $CH_3CN$  in the APCI source of a commercial mass spectrometer LTQ XL (Thermo Scientific) [2,3] and reacted with  $HC_3N$  in the trap to produce heavier anions through the  $CN^- + x HC_3N \rightarrow (HC_3N)_y C_{2p+1}N^- + z HCN$  processes. These product anions are then mass-selected in the trap and irradiated with VUV photons (5-21 eV) from the DESIRS beamline. Their decay is followed as a function of irradiation time as illustrated in Figure 1.




Figure 1: Trap ion induced loss by photodetachment: comparison of the exponential decay of the measured  $(HC_3N)_3C_5N^-$  anion signal as a function of the irradiation time with and without synchrotron light for a photon energy of 8 eV.

## Acknowledgements

Programme National de Planétologie (PNP), COST (Action CM0805 « The Chemical Cosmos »), Czech Science Foundation (Grant No. P208/11/0446), (Grant Nos. OC10046).

## References

- [1] J. Zabka, C. Romanzin, C. Alcaraz, M. Polasek, *Icarus*, 219, pp. 161-167, 2012.
- [2] A. R. Milosavljevic, C. Nicolas, J. Lemaire, C. Dehon, R. Thissen, J.-M. Bizau, M. Refregiers, L. Nahon and A. Giuliani, *Phys.Chem.Chem.Phys.*, 13, pp. 15432-15436, 2011.
- [3] A. R. Milosavljevic, C. Nicolas, J.-F. Gil, F. Canon, M. Refregiers, L. Nahon and A. Giuliani *J. Synchrotron Rad.* 19, pp. 174-178, 2012.