

Discovery of the double impact crater in the Imilchil region, Morocco

A. Ibhi (1), H. Nachit (1), El H. Abia (1), A. Ait Touchnt (2) and C. Vaccaro (2)

(1) Geoheritage and Geomaterials Laboratory, University of Ibn Zohr, B.P. 8106, Morocco. (2) Department of Earth Sciences, P1-SZA103 Via Saragat, 1. University of Ferrara, Ferrara-44100, Italy. (a.ibhi@uiz.ac.ma / Fax : +212-528220100)

Abstract

Isli and Tislit, the Moroccan "Romeo and Juliette" according to a popular legend gave their name to a twin lakes in Imilchil. There we show evidences that these two lakes or impact craters. The proof as follow: (i) the fragments of iron meteorites found on both sides of Isli and Tislit lakes belong to the same parent meteorite and siderite type, (ii) The sedimentary formations which are almost tabular in the area of the lakes become tilted with a centripetal sloping all around and towards Isli lake, (iii) the circular shape of the Tislit lake is similar to that of the Isli lake, (iv) the impact breccias found on the south of the Isli lake contain shocked quartz crystals (v) the radial fractures of high pressure affect some sedimentary beds. All these observations give evidence of the presence of an impact crater due to the fall of meteorites.

1. Introduction

More than 183 structures of meteoritical impacts are highlighted on the Earth [1], including only three dual impact craters reported [2]. The impact craters of Isli and Tislit, 32°13'N, 05°33'W; 32°12'N, 05°38'W are respectively, located at 10 km and 4 km from the Imilchil village on the Central High Atlas in Morocco at respective altitudes of 2272 and 2266 meters. They are spaced apart from each other by 9.4 km. The origin and the date of formation of these two lakes are still unknown today. Surveys undertaken near them and in the surrounding areas leads to believe that they correspond to a dual meteoritical impact crater. We present here the discovery of this new dual impact crater in the Central High Atlas in Morocco. It is interesting, first by its young age, its perfect conservation and the geomorphology of the two depressions (Figures 1).



Figure 1. Panorama of the Isli Lake.

2. New observations

2.1 Meteorite

The first fragment of this meteorite was harvested by a nomad in June 2012. The members of the laboratory harvested many fragments of this meteorite well preserved in the quaternary (Upper Soltanian) layers, dated 40,000 years BP [3]. The collected fragments on the both sides of the both impact craters and those collected on the surrounding plains are of the same nature, thus having fragments produced by a large explosion by the impactor during a collision with the soil's surface. Samples weighing 100 g to more than 9 kg (Figures 2A, 2B and 2C) were harvested at several kilometers from the west to the east of the craters.

The samples found have a rusty appearance, with scales of alteration of several centimeters thick of oxides and iron hydroxide covering the entire outer surfaces thereof, indicating a long residence time in the soil. The polished surfaces don't show the Widmanstätten figures. The meteorite of Imilchil is an Ataxite rich in nickel, it consists mainly of iron mixed with 17, 3% nickel (X-ray fluorescence method in the department of Earth sciences, the University of Ferrara). The scanning electron microscopy revealed the presence of thin diffuse bands of kamacite (around 25 µm wide) in a matrix of plessite (Figure 2D) this texture is the evidence of structural deformation due to shock impact.



Figure 2. Meteorite fragments of Imilchil. A: Sample of more than 9 kg. B: Sample of more than 1.5 kg. C: Sample of 290 grams on site. D: Band of kamacite in a plessite matrix.

2.2 Impact craters

The dual impact craters of Imilchil (Isli and Tislit) affect the jurassic sedimentary formations (red layers). The both craters are practically circular and they have respectively 1500 and

1000 meters in diameter. The crater Isli is an enough regular bowl with very pronounced slopes; its maximum depth is 95 meters, the seismic profiles obtained reveal that the thickness of sediments filling the lake is 100 meters; the oldest have an age of about 35,000 years [4]. The sedimentary formations that are almost tabular in the platform lakes become inclined with centripetal dips, all around, towards the Isli lake forming thus its watershed.

Polyimict and monomict breccias were observed at the edges of these craters (Figures 3A and 3B), they don't form big masses, the erosion was probably erased most. They consist of irregular fragments with millimeter and centimeter sizes, in a carbonate matrix. In the breccias collected on the southern edges of the Isli crater, we noted the presence of several shocked quartz grains in planar structures (FP) [5-6] associated sometimes with rolling extinction, indicators of shock metamorphism (Figure 3C). The most quartz grains have a single family of PFs, but some grains have two families of planar structures. These breccias could be the result of debris ejecta. We could also observe radial fractures (Figure 3D) which could have been produced by impacts [7].




Figure 3. Metamorphism impact figures. A: polymict breccias. B: monomict breccias. C: shocked quartz showing PFs. D: radial fractures.

3. Discussion and conclusion

Our observations lead to a structural outline of an impact. The fragments of meteorites (around 180 Kg) found in the area all belong to one and the same parent meteorite called Ataxite. Sedimentary layers are verticalized around the Isli crater; outside the edges of the crater, there is no evidence of vertical layers. This is most likely due to a meteorite fall [8]. Witnesses of impact breccias are still present on the edges of craters, although most of these breccias had been erased by

erosion. These breccias contain shocked quartz grains with planar structures indicators of shock metamorphism [5-6].

In conclusion, based on these observations we can confirm that the circular structures of Isli and Tislit discovered in the north of Imilchil (Morocco) were produced by the impact of a small asteroid with more than 100 meters in diameter.

The dual impact crater, oriented est-west, resulted from the fragmentation of a meteorite at its input in the Earth's atmosphere in two pieces which then produced two craters, oriented along the trajectory of the fireball [9-10]. This phenomenon has been observed in three double craters identified in the world [2]. According to the state of conservation of the meteorite into the sedimentary layers "Soltanian" and the dating of ancient lake sediments of the Isli lake [4], we can estimate their age limit to 40,000 years.

References

- [1] J. Spray, "Earth Impact Database," 2011. <http://www.passc.net/EarthImpactDatabase/index.html>
- [2] J. N. Salomon and T. Auly, "Impactors and astroblems: attempt of classification and geographical approach," *Géomorphologie*, 2010, pp. 3-20. ISBN 2-913282-47-4.
- [3] A. Weisrock and M. Fontugne, "Morphogenèse éolienne littorale au Pléistocène supérieur et à l'Holocène dans l'Oulja atlantique marocaine," *Quaternaire*, Vol. 3, 1991, pp. 164-175.
- [4] E. Zeroual, "Enregistrements climatiques dans les sédiments de lac Isli (Haut Atlas, Maroc)," Ph.D. Thesis, Cadi Ayyad University, Morocco, 1995.
- [5] D. Stöffler and F. Langenhorst, "Shock metamorphism of quartz in nature and experiment : I. Basic observations and theory," *Meteoritics & Planetary Science*, Vol. 29, 1994, pp. 155-181.
- [6] R. A. F. Grieve, F. Langenhorst and D. Stöffler, "Shock metamorphism of quartz in nature and experiment," *Meteoritics & Planetary Science*, Vol. 31, 1996, pp. 6-35.
- [7] A. Sagy, Z. Reches and J. Fineberg, "Dynamic fracture by large extraterrestrial impacts as the origin of shatter cones," *Nature*, Vol. 418, 2002, pp. 310-313.
- [8] J. F. Moyen, "La structure d'impact de Vredefort," In M. J. Van Kranendonk, R. H. Smithies and V. Bennett, Ed., *Earth's Oldest rocks*, Elsevier, 2007, pp. 669-698. http://www.academia.edu/1107614/La_structure_dimpact_de_Vredefort
- [9] M. R. Dence, "Meteorite breakup," *Nature*, Vol. 289, No 198, pp. 346-351.
- [10] B. M. French and C. Koeberl, "The convincing identification of terrestrial meteorite impact structures: What works, what doesn't, and why," *Earth-Science Reviews*, Vol. 98, 2010, pp. 123-170.