

Fluvial processes in eastern Hellas Planitia, Mars – Insights from age determinations

W. Zuschneid, S. van Gasselt

Freie Universität Berlin, Institute of Geological Sciences, Planetary Sciences and Remote Sensing Group,
Malteserstr. 74-100, 12249 Berlin (wilhelm.zuschneid@fu-berlin.de)

1. Introduction

With a diameter of 2,300 km and a depth of more than 8,000 m, the Hellas basin is one of the major geomorphic and topographic features in the southern Martian hemisphere. It has been acting as a depositional sink since its formation 4 Ga ago [1] and has been the location of a wide variety of geologic processes. In this ongoing study, we investigated the stratigraphy of fluvial features in the eastern Hellas region on Mars, focusing on apparently young fluvial depositional areas in the Dao and Harmakhis Valles region and on the eastern plains within the Hellas basin.

The region connecting Hesperia Planum with Hellas Planitia is characterized by a patchwork of remnant massifs protruding through a set of plains units of varying morphology and surface texture, indicating a diverse and eventful geologic history. A variety of geologic processes has been observed, from the basin-forming Hellas impact, widespread plains-forming and central vent volcanism [2,3] to fluvial and cold climate processes [3,4]. In order to analyze the stratigraphy, we performed crater counts for selected areas to obtain absolute ages for young areas formed by fluvial processes. We also obtained a lower limit for the age of major fluvial processes.

2. Data and Methods

The study is based on the analysis of image data collected by the HRSC (Mars Express), the CTX camera (MRO) and the THEMIS IR instrument (Mars Odyssey). CTX data was processed and map-projected using the ISIS suite, and then integrated into ArcGIS for further analysis and supplemented by the THEMIS IR daytime mosaic produced by ASU. Areas for crater counting and age determination were selected based on the apparent youth of the surface, and their potential association with fluvial landforms.

In addition, several additional areas with different origin were selected for comparison and to complement the general picture. Problems arose from a reduced image quality due to atmospheric dust at lower elevations in the Hellas basin.

Using the CraterTools 2.2 extension [5], counting areas were outlined based on the criteria as outlined e.g. in [6], care was taken to exclude obvious clusters and chains of secondary craters. The crater counts obtained were analyzed using the Craterstats 2 software described in [7].




Fig 1.: Eastern Hellas basin: overview of crater counting areas (red outline) on color-coded MOLA shaded relief. Scale bar length is 750 km.

3. Observations

The eastern region of Hellas Planitia and the adjacent rim region are characterized by abundant fluvial features [8]. The most obvious are Dao and Harmakhis Valles, but plenty of smaller fluvial channels can be observed at lower elevations along the slope into the basin. Associated with these smaller channels are flat, uniform small plains, which are interpreted as floodplains. At the channel termini on the eastern plains in Hellas basin, associated sedimentary deposits are observed.

We investigated selected areas of this type along a small valley north of Dao Vallis, and along Sungari Vallis, a medium-sized channel located between Dao and Harmakhis Valles; in addition, we investigated the deposits at the termini of the valleys. For these areas, crater counts were performed to obtain crater size-frequency distributions and ages. In addition, craters were counted in several of the shallow basins separated by ridges located on the eastern floor of the Hellas basin [9]. This large spatial spread of counting areas made an evaluation of the spatial extent of resurfacing effects possible.




Fig. 3: Age distribution of measured ages in selected counting areas. Ages between 3.2 Ga and 3.75 Ga represent the formation of the basin floor units, younger ages represent the subsequent modification. The abundance of ages between 0.75 Ga and 0.25 Ga indicates a widespread episode of resurfacing of the eastern Hellas ridged plains.

3. Results

Although the crater count areas are spatially separated, the absolute age determinations yield relatively consistent results, indicating ages for the last fluvial resurfacing between 250 and 750 Ma. Similar resurfacing ages were obtained for the basins on the eastern plains. Pre-resurfacing ages of 3.4 to 3.7 Ga were determined for the basins, establishing a baseline for the oldest events following the formation of Hellas basin. The formation ages of > 3.5 Ga for the plains are in good agreement with results obtained by [2,6] (for much larger counting areas).

3. Conclusions

The results of our crater counts indicate an episode of increased fluvial activity from 250 to 750 Ma. This

fluvial activity led to the partial resurfacing of areas on the slope as well as on the basin floor. On the slope into the Hellas basin, blanketing of surfaces adjacent to fluvial valleys suggests that resurfacing occurred by sedimentary deposition. The extent of the depositional areas indicates a high sediment load associated with the single fluvial events. The water sources associated with the possible fluvial activity were not in the focus of this investigation, but a release from the subsurface (shallow or deep) has been proposed for Sungari Vallis, e.g. by [10, 11] and also seems feasible for the other, smaller valley system north of Dao Vallis.

For the basins on the Hellas floor, a resurfacing by sedimentary deposition is not directly observed. Here, better HiRISE coverage may be helpful in the identification of resurfacing mechanisms. Crater counts on an adjacent remnant massif indicate that not all surfaces were affected by the resurfacing episode. Older resurfacing events were also identified (e.g. at around 1.9 Ga), but their occurrence is not as evident in the crater counts as the younger episode.

Acknowledgements

This work was supported by the DLR Space Administration on behalf of the Federal Ministry for Economic Affairs and Energy, grant 50QM1301 (HRSC on Mars Express). We thank the HRSC experiment team at DLR Berlin and the HRSC operations team at ESOC for their successful planning, acquisition and processing of the HRSC data.

References:

- [1] Werner, S. (2008) Icarus, 195, 45-60, [2] Williams, D.A. et al. (2010) E&PSL 294 3-4, 492-505, [3] Crown, D. A. et al. (1992) Icarus, 100, 1-25, [4] van Gasselt, S. et al. (2007) JGR 112 E9, [5] Kneissl et al. (2011) P&SS 59, 11-12, 1243-1254, [6] Platz, T., et al. (2013) Icarus, 225, 806-827, [7] Michael, G. G., and Neukum, G. (2010) E&PSL 294 3-4, 223-229, [8] Bleamaster, L. F., and Crown, D. A. (2010) USGS Sci. Inv. Maps, I-3096 [9] Zuschneid, W., and van Gasselt, S. (2013) LPSC XXXIV, Abstract #2191, [10] Squyres, S. W. et al. (1987) Icarus 70, 385-408, [11] Kostama, V.-P. et al. (2010) E&PSL 294 3-4, 321-331