

## The challenge and scientific application of the CO<sub>2</sub> 4.3 $\mu$ m atmospheric limb emission of Mars

M. A. López-Valverde (1), A. Piccialli (2), M. Giuranna (3), B. Funke (1), M. López-Puertas (1), A. A. Jurado-Navarro (1), M. García-Comas (1), F. González-Galindo (1), J. J. López-Moreno (1) and S. Jiménez-Monferrer (1)

(1) Instituto de Astrofísica de Andalucía/CSIC, Granada, Spain. (valverde@iaa.es / Fax: +34-958-814530)

(2) LESIA, Observatoire de Paris, Meudon, France

(3) Istituto di Astrofisica e Planetologia Spaziali/INAF, Rome, Italy

### Abstract

The atmospheric fluorescent emissions of CO<sub>2</sub> at 4.3- $\mu$ m have been observed in the daytime upper atmosphere of Mars from a limb geometry by the instruments OMEGA and PFS on board Mars Express [1, 8]. Initial analysis using non-local thermodynamic equilibrium (NLTE) models show that the emissions are well understood [7, 3, 6]. Yet they have not been exploited to derive important thermospheric parameters, like CO<sub>2</sub> densities and temperatures. Our major goals are to improve current NLTE models with a joint study of OMEGA and PFS data, and to build an ambitious state-of-the-art NLTE retrieval scheme for Mars. Recent progress has been made in these directions on Mars, Venus and Earth. We will present a summary of these efforts and the difficulties and expectatives for its application to the Mars Express data.

### 1. Introduction

Infrared NLTE emissions by atmospheric molecules, specially under fluorescent or solar-pumping situations, are powerful tools to sound the upper layers of a planetary atmosphere [5]. Limb observations, in addition, are specially suited for a proper vertical sounding. The particular emissions by the ro-vibrational bands of CO<sub>2</sub> around 4.3  $\mu$ m share these two advantages and are among the strongest molecular emissions in the infrared in the three terrestrial planets [6]. The instruments OMEGA and PFS have been performing systematic observations of these emissions on board Mars Express since 2003 and form a dataset which needs to be properly exploited. However, their scientific exploitation has a couple of difficulties, inherent to the NLTE nature of the emissions. One of them is the need to develop a suitable NLTE forward model (based on the theoretical NLTE population models) which produce simulated radiances in suffi-

ciently good agreement with the observations. Another difficulty is linked to the inherent uncertainties in the microscopic description of the NLTE populations involved in a CO<sub>2</sub> atmosphere like that of Mars. Specifically, the lack of laboratory information on collisional energy exchanges between CO<sub>2</sub> states at high vibrational states and with isotopes. Actually, when spectral resolution and sensitivity are good enough, insight into and retrieval of some collisional rate coefficients are possible from remote sounding [4]. A third difficulty is the large optical thickness of these emissions in a limb geometry, which may produce variations in the actual location of the “emitting region” along the line-of-sight. Retrievals under optically thin conditions, free from this last difficulty because the emitting region is at the tangent point, have recently been carried out for CO at 4.7  $\mu$ m in the upper atmosphere of Venus [2]. A careful evaluation of all these conditions and difficulties, and a proper fit of the OMEGA and PFS data with a NLTE forward model are needed in order to evaluate the ability to investigate the upper Martian atmosphere using these datasets, and are the focus of an ongoing effort in our teams.

### 2. Recent NLTE results

#### 2.1. Mars upper atmosphere

The OMEGA and PFS datasets seem to obey the expected NLTE behaviour of the emissions regarding the two most important aspects: spectral shape and absolute magnitude of the emitted radiance. Also the variation of these with altitude and solar illumination seem to be understood, at least in a global or averaged sense [7, 3]. In order to explain the spectral shape measured by OMEGA, however, changes in vibrational-vibrational exchanges between CO<sub>2</sub>( $\nu_3$ ) states was needed; a solution which also improved similar measurements by VIRTIS/Vex in the Venus thermosphere

[6]. An extended study using both OMEGA and PFS is needed, in particular the PFS higher spectral resolution, can be very useful to settle this question.

## 2.2. Venus upper atmosphere

In the Venus upper atmosphere, the 4.3  $\mu\text{m}$  emissions by  $\text{CO}_2$  present similar difficulties. A simpler retrieval (optically thin situation) was applied to the limb fluorescent emissions by CO at 4.7  $\mu$  from VIRTIS/Venus Express, obtaining both CO abundance and temperatures simultaneously [2]. CO observations in Mars, at least available from PFS, should also be exploited, and a joint retrieval with  $\text{CO}_2$  at 4.3  $\mu\text{m}$  shall be particularly useful to obtain the total carbon content in the Martian thermosphere.

## 2.3. Earth upper atmosphere

An interesting study in the Earth's upper atmosphere using MIPAS/Envisat observations has been recently conducted in the Granada group [4]. Jurado-Navarro and colleagues not only derived  $\text{CO}_2$  but also rate coefficients of a few vibrational-to-vibrational and vibrational-to-translation energy transfer routes involving  $\text{CO}_2$  states. Unfortunately the OMEGA and PFS spectral resolutions are not as good as those of MIPAS, but the NLTE retrieval scheme used by these authors should be applied to the Martian case to examine the possibility of similar derivations. In the case of Mars, or Venus, the collisional exchanges driving the populations are slightly different, and offer new insight into the governing NLTE mechanisms in a  $\text{CO}_2$  atmosphere.

## 3. Summary and Conclusions

The NLTE limb emissions of  $\text{CO}_2$  at 4.3  $\mu\text{m}$  observed by Mars Express are a unique dataset to extend our description of the upper mesosphere and the thermosphere of Mars. The understanding of the emissions, their correct simulation and the NLTE retrieval of densities and temperatures are our major objectives. A further strategy behind this work is the preparation of suitable retrieval tools for the analysis of the upcoming ExoMars Trace Gas Orbiter (EMTGO), specifically to exploit infrared measurements of the upper atmosphere subject to NLTE features in a completely different observing mode (atmospheric absorption) from two of the EMTGO instruments: NOMAD and ACS.

## Acknowledgements

This work has received funding from the European Union's Horizon 2020 Programme (H2020-Compet-08-2014) under grant agreement UPWARDS-633127

## References

- [1] V. Formisano, A. Maturilli, M. Giuranna, E. D'Aversa, and M. A. López-Valverde. Observations of non-lte emission at 4.5 microns with the planetary fourier spectrometer aboard the mars express mission. *Icarus*, 182:51–67, May 2006.
- [2] G. Gilli, M. López-Valverde, J. Peralta, S. Bougher, A. Brecht, P. Drossart, and G. Piccioni. Carbon monoxide and temperature in the upper atmosphere of venus from virtis/venus express non-lte limb measurements. *Icarus*, 248(0):478 – 498, 2015.
- [3] M. Giuranna, A. Kutepov, L. Rezac, A. Feofilov, and V. Formisano. Analysis and modelling of PFS/MEx limb observations of 4.3- $\mu\text{m}$   $\text{CO}_2$  non-LTE emission. In *European Planetary Science Congress 2012*, page 682, Sept. 2012.
- [4] A. A. Jurado-Navarro, M. Lopez-Puertas, B. Funke, M. García-Comas, A. Gardini, G. P. Stiller, and T. von Clarmann. Vibration-vibration and vibration-thermal energy transfers of  $\text{CO}_2$  with  $\text{N}_2$  from mipas high resolution limb spectra. *J. Geophys. Res.*, submitted, 2015.
- [5] M. López-Puertas and F. W. Taylor. *Non-LTE radiative transfer in the Atmosphere*. World Scientific Pub., Singapore, 2001.
- [6] M. A. López-Valverde, M. López-Puertas, B. Funke, G. Gilli, M. García-Comas, P. Drossart, G. Piccioni, and V. Formisano. Modelling the Atmospheric Limb Emission of  $\text{CO}_2$  at 4.3  $\mu\text{m}$  in the Terrestrial Planets. *Planet. Space Sci.*, 59:988–998, 2011.
- [7] A. Piccialli, P. Drossart, M. A. Lopez-Valverde, F. Altieri, A. Määttänen, B. Gondet, O. Witasse, and J. P. Bibring. Characterization of OMEGA/MEx  $\text{CO}_2$  non-LTE limb observations on the dayside of Mars. In *European Planetary Science Congress 2012*, page 504, Sept. 2012.
- [8] A. Piccialli, E. Palomba, M. A. López Valverde, G. Gilli, D. Grassi, M. D'Amore, and V. Formisano. Pfs-mex nadir observations at 4.3  $\mu\text{m}$  in the mars atmosphere with account for non-lte emissions. In *European Planetary Science Congress 2006*, pages 257–+, 2006.