

Dynamically interesting asteroids discovered at the VATT Observatory

I. Włodarczyk (1), K. Černis (2) and R. Boyle (3)

(1) Chorzow Astronomical Observatory (IAU 553), Al. Planetarium 4, Chorzow 41-500, Poland (e-mail:astrobit@ka.onet.pl), (2) Institute of Theoretical Physics and Astronomy, Vilnius University, Goštauto 12, Vilnius LT-01108, Lithuania (e-mail:Kazimieras.Cernis@tfai.vu.lt), (3) Vatican Observatory Research Group, Steward Observatory, Tucson, Arizona 85721, U.S.A. (e-mail:rpboyle@mac.com)

Abstract

We present results of a long-term research program on dynamically interesting asteroids discovered at the VATT Observatory.

1. Introduction

One of the research projects of the Vatican Observatory is a study of asteroids for a better prediction of their orbits. The observing program on the 1.8 m telescope VATT is targeted on observing the Main Belt and the Centaur group asteroids, Trans-neptunian objects (TNO), as well as on the search of new objects. First astrometric observations of asteroids and comets with the VATT at the Mt. Graham Observatory (IAU code 290, longitude 109.89201 W, latitude 32.70133 N, altitude 3178 m) were initiated in 2000-2001 using CCD images exposed in the focal plane of the VATT [1].

The first new asteroid, (220696) = 2000 YJ143, was discovered in 2000 by W. H. Ryan. In 2007, 75 new asteroids were discovered by C. W. Hergenrother. A new observational program for the search and observations of asteroids was started in 2009 by R. P. Boyle and K. Černis.

2. Orbits

We continue presentation of dynamically interesting asteroids discovered at the VATT Observatory. They are described in:

- Centaur Asteroid 2012 DS85 [5]
- Amor-class asteroid 2012 XH16 [6]
- Transneptunian (TNO) object (420356) 2012 BX85 Praamžius [7]
- Distant Object (463368) 2012 VU85 [8]

Fig. 1 presents orbits of four discovered asteroids at VATT Observatory in the ecliptic plane: 2012

DS85, 2012 XH16, (420356) 2012 BX85 Praamžius and (463368) 2012 VU85.

The first object observed was an asteroid of the Centaur group 2009 HW77 [2],[4] in April of 2009. All published orbits of asteroids discovered at the VATT Observatory were computed by I. Włodarczyk using the OrbFit software (<http://adams.dm.unipi.it/~orbmaint/orbfit/>). Accuracy of computations is described in [3].

3. Summary and Conclusions

We presented short history of asteroids discovered at the VATT Observatory. Starting orbits of four selected asteroids are presented in Fig. 1.

References

- [1] Hergenrother C.W. and Spahr T.B. 2001, MPC 42751
- [2] Černis K., Eglitis I., 2009, MPEC, 2009-U68
- [3] Włodarczyk I., 2009, Icar, 203, 119
- [4] Włodarczyk I., Černis K., Eglitis I., 2011, MNRAS, 418, 2330
- [5] Černis K., Boyle R. P., Laugalys V., Włodarczyk I., 2012, BaltA, 21, 455
- [6] Włodarczyk I., Černis K., Boyle R. P., Laugalys V., 2014, MNRAS, 438, 2621
- [7] Černis K., Boyle R. P., Włodarczyk I., 2016, BaltA, 25, 189
- [8] Włodarczyk I., Černis K., Boyle R. P., 2017, AcA, 67, 81




Figure 1: Orbits of selected asteroids discovered at VATT Observatory presented in the ecliptic plane. The dashed line denotes the part of the orbit below the ecliptic plane.