

Imaging of energetic neutral atoms with the Jovian Neutral Atoms Analyser onboard JUICE: Charge exchange ENAs near Ganymede

M.B. Neuland (1), Y. Futaana (1), S. Fatemi (1), M. Shimoyama (1), K. Asamura (2), A. Vorburger (3), P. Wurz (3), M. Wieser (1) and S. Barabash (1)

(1) Institutet för rymdfysik, IRF, Rymdcampus 1, SE - 98128 Kiruna, Sweden

(2) Institute of Space and Astronautical Science, ISAS, Japan Aerospace Exploration Agency (JAXA)
Yoshinodai 3-1-1, Chuo-ku, Sagami-hara, Kanagawa, 252-5210, Japan

(3) University of Bern, Physics Institute, Space Research and Planetary Sciences, Sidlerstrasse 5,
CH - 3012 Bern, Switzerland

Abstract

The Jovian Neutral Atoms Analyser (JNA) is one of the sensors of the Particle Environment Package (PEP), which is one instrument suite of the JUICE scientific payload. Mapping of the energetic neutral atoms (ENAs) in the Ganymede magnetosphere, will offer valuable clues about the interaction of the Jovian plasma with the icy moon's surface and the magnetospheric plasma processes. We will present expectations of neutral fluxes, particularly neutrals produced via charge exchange, in the vicinity of Ganymede. From these expectations from our model we infer the expectations for JNA measurements.

1. Introduction

The Jovian system is subject to complex plasma physics processes and can therefore be considered as a giant particle accelerator. Due to the plasma interactions, the Jovian moons are affected by diverse processes that lead to a vast range of space weathering effects and to a constant mass input to the environment of the moons.

Ganymede possesses a strong intrinsic magnetic field that imposes restrictions on the possible trajectories of plasma ions in the Ganymede magnetosphere [1]. As a result, certain terrains on the surface of the Jovian moon are protected against space weathering processes while others are not.

The JNA onboard the JUICE spacecraft will measure energetic neutral light and heavy atoms in an energy range from 10eV to 3keV [2], which covers the energy range of ENAs emanating from various different ENA production mechanisms. Low energy ENAs produced via sputtering and backscattering

will be used to image the precipitation regions and, in particular, a boundary of open and closed field lines. In addition, ENAs are expected to be produced by the charge – exchange mechanism in the vicinity of Ganymede. From the measurement of these ENAs, we can infer the global plasma distribution in the Ganymede magnetosphere.

In this study we investigate the interaction between the Jovian plasma and the Ganymede magnetosphere and exosphere. In the Jovian system these interactions of the co-rotating magnetospheric plasma with the icy moons give rise to several observable effects, like for example the UV-aurorae observed on Ganymede [6].

The energy balance in terms of mass and radiation balance is one key scientific question of the mission as well as the surface and exosphere composition of the Jovian moons [4]. The direct measurements of mass flux by the JNA will undoubtedly contribute to answer these questions.

2. Model

To calculate the expected fluxes of ENAs as well as the number of ENAs from the charge exchange processes, we combined the plasma spatial distribution modelled by hybrid simulation [5], the exospheric density model for Ganymede [8], and the charge exchange cross sections [7]. Using a hybrid model, we derived the fraction of ENAs from charge exchange processes and the expected neutral fluxes in the JNA instrument.

Fig. 1 shows the plasma ion locations from [5] in the GPhiO system as a snapshot in time. In the graph, ions are shown in white colour, thus bright regions

designate higher plasma density. The co-rotating Jovian plasma precipitates on the Ganymede magnetosphere from the rear, causing a void in front of the moon, appearing in black in Fig.1, seen in direction of its trajectory along the orbit.




Figure 1: Plasma spatial distribution in the vicinity of Ganymede, reproduced from [5].

3. Summary

The JNA instrument, one of the sensors of the PEP scientific payload on the JUICE mission, will measure ENAs in the Ganymede magnetosphere and exosphere. These measurements will greatly contribute to our understanding of the interaction of the Jovian plasma with Ganymede, its icy surface and its intrinsic magnetic field. Based on a hybrid model we derived the neutral fluxes produced by charge exchange processes of high energetic plasma particles and neutral atoms in the Ganymede exosphere and the resulting fluxes expected to be measured by the JNA. We present our model, the resulting observations for species heavier than N and the ENA fluxes along the JUICE Ganymede orbit expected to be measured by the JNA instrument.

Acknowledgements

This work is supported by the Swiss National Science Foundation (SNF) Early PostDoc Mobility Grant No.168708 and the Swedish National Space Board (SNSB).

References

- [1] M.G. Kivelson, K.K. Khurana and M. Volwerk, The Permanent and Inductive Magnetic Moments of Ganymede, *Icarus* 157, 2002.
- [2] O. Grasset et al., Jupiter Icy moons Explorer (JUICE): An ESA mission to orbit Ganymede and to characterise the Jupiter system, *Planet. Space Sci.* 78, 2013.
- [3] M. Wieser et al., Emission of energetic neutral atoms from water ice under Ganymede surface-like conditions, *Icarus* 269, 2016.
- [4] J.J. Plaut et al., Jupiter Icy moons Explorer (JUICE): Science objectives, mission and instruments, 45th Lunar and Planetary Science Conference 2717, 2014.
- [5] Fatemi et al., On the formation of Ganymede's surface brightness asymmetries: Kinetic simulations of Ganymede's magnetosphere, *Geophys. Res. Lett.* 43(10), 2016.
- [6] P.D. Feldman et al., HST/STIS Ultraviolet Imaging of Polar Aurora on Ganymede, *Astrophys. J.* 535 (2), 2000.
- [7] Scherer et al., Ionization rates in the heliosheath and in astrosheaths - Spatial dependence and dynamical relevance, *Astronomy & Astrophysics* 563, 2014.
- [8] P. Wurz et al., The Exospheres of Europa, Ganymede and Callisto, Conference paper, *Exo - Climes III*, Bern/Switzerland, March 2014