

Model expectations for the D/H distribution on Mars as observed by NOMAD

Frank Daerden (1), Lori Neary (1), Sébastien Viscardy (1), Shohei Aoki (1,8,9), Arianna Picciali (1), Séverine Robert (1), Valérie Wilquet (1), Ian B. Thomas (1), Bojan Ristic (1), Ann Carine Vandaele (1), Geronimo Villanueva (2), Michael J. Mumma (2), Robert E. Novak (3), Thérèse Encrenaz (4), Stephen Lewis (5), James Holmes (5), José Juan Lopez-Moreno (6), Giancarlo Bellucci (7), Manish Patel (6), and the NOMAD team.

(1) Royal Belgian Institute for Space Aeronomy BIRA-IASB, Brussels, Belgium, (2) NASA Goddard Space Flight Center, Greenbelt, MD, USA, (3) Iona College, New Rochelle NY, USA, (4) LESIA, Observatoire de Paris, CNRS, UPMC, UPD, Meudon, France, (5) Open University, Milton Keynes, UK, (6) Instituto de Astrofísica de Andalucía (IAA-CSIC), Granada, Spain, (7) Instituto di Astrofisica e Planetologia Spaziali (IAPS), Istituto Nazionale di Astrofisica (INAF), Rome, Italy, (8) Fonds National de la Recherche Scientifique, Belgium, (9) Tohoku University, Japan (Frank.Daerden@aeronomie.be)

Abstract

The NOMAD (“Nadir and Occultation for Mars Discovery”) spectrometer suite on board the ExoMars Trace Gas Orbiter (TGO) has been designed to investigate the composition of Mars’ atmosphere, with a particular focus on trace gases, clouds and dust, and started its science operations in April 2018. One of the primary science objectives of NOMAD is the detailed investigation of semiheavy water (HDO), both in nadir (total columns) and in solar occultation (vertical profiles). Interestingly, NOMAD has the capability to simultaneously retrieve H₂O and HDO, hence the very important D/H ratio. This ratio provides information on cloud formation and atmospheric escape. The GEM-Mars model was recently extended with HDO fractionation. We present some first results that will be useful to guide the interpretation of D/H data from NOMAD.

Introduction

The isotopic composition of fresh water on Earth has been defined by the Vienna Standard Mean Ocean Water (VSMOW), in which the HDO isotopologue of water has a relative abundance (D/H or $\frac{1}{2} \times \text{HDO}/\text{H}_2\text{O}$) of 155.76 ± 0.1 ppm (parts per million). In the atmosphere of Mars, D/H has been found to be enhanced by 5-7 times compared to Earth [1, 2 and references therein], indicating a stronger atmospheric depletion of H₂O on Mars compared to Earth. Until now, HDO and D/H have been observed from Earth, and NOMAD will be the first instrument in orbit to provide detailed measurements. D/H fractionation is expected to occur in the Martian atmosphere following two processes: upon condensation/

evaporation of water, where the heavier isotopologue will condense/vaporize at slightly higher temperature, and by photolysis, where HDO has a lower absorption cross-section in the UV [3]. Also, the heavier isotope will escape less readily above the exobase during Jeans escape. Also the D/H ratio in ancient water ice reservoirs, such as the permanent water ice cap, is expected to impact on the atmospheric value [2].

Modeling the D/H ratio on Mars

The GEM-Mars General Circulation Model [4] includes both the formation of water ice clouds and photochemistry. As such, it is well equipped to do simulations of the D/H fractionation. First results will be presented and interesting features will be highlighted. Model expectations will be discussed in view of the observational geometries of NOMAD and if possible compared to first results (which will be presented in Aoki et al., this session). For example, the model can help to understand how representative observed D/H ratios are in the solar occultation mode, where measurements are confined to the terminator and could be affected by morning/evening cloud and fog formation.

Acknowledgements

The NOMAD experiment is led by the Royal Belgian Institute for Space Aeronomy (IASB-BIRA), assisted by Co-PI teams from Spain (IAA-CSIC), Italy (INAF-IAPS), and the United Kingdom (Open University). This project acknowledges funding by the Belgian Science Policy Office (BELSPO), with the financial and contractual coordination by the ESA

Prodex Office (PEA 4000103401, 4000121493), by the Spanish Ministry of Economy, Industry and Competitiveness and by FEDER funds under grant ESP2015-65064-C2-1-P (MINECO/FEDER), by the Italian Space Agency through grant 2018-2-HH.0, as well as by UK Space Agency through grant ST/P000886/1. US investigators were supported by the National Aeronautics and Space Administration. The research was performed as part of the “Excellence of Science” project “Evolution and Tracers of Habitability on Mars and the Earth” (FNRS 30442502).

The NOMAD Team

Scientific team: Vandaele, Ann Carine; Lopez Moreno, Jose Juan; Bellucci, Giancarlo; Patel, Manish; Allen, Mark; Alonso-Rodrigo, Gustavo; Altieri, Francesca; Aoki, Shohei; Bauduin, Sophie; Bolsée, David; Clancy, Todd; Cloutis, Edward; Daerden, Frank; D'Aversa, Emiliano; Depiesse, Cédric; Erwin, Justin; Fedorova, Anna; Formisano, Vittorio; Funke, Bernd; Fussen, Didier; Garcia-Comas, Maia; Geminale, Anna; Gérard, Jean-Claude; Gillotay, Didier; Giuranna, Marco; Gonzalez-Galindo, Francisco; Hewson, Will; Homes, James; Ignatiev, Nicolai; Kaminski, Jacek; Karatekin, Ozgur; Kasaba, Yasumasa; Lanciano, Orietta; Lefèvre, Franck; Lewis, Stephen; López-Puertas, Manuel; López-Valverde, Miguel; Mahieux, Arnaud; Mason, Jon; Mc Connell, Jack; Mumma, Mike; Nakagawa, Hiromu; Neary, Lori; Neefs, Eddy; Novak, R.; Oliva, Fabrizio; Piccialli, Arianna; Renotte, Etienne; Robert, Severine; Sindoni, Giuseppe; Smith, Mike; Stiepen, Arnaud; Thomas, Ian; Trokhimovskiy, Alexander; Vander Auwera, Jean; Villanueva, Geronimo; Viscardy, Sébastien; Whiteway, Jim; Willame, Yannick; Wilquet, Valérie; Wolff, Michael; Wolkenberg, Paulina – *Tech team:* Alonso-Rodrigo, Gustavo; Aparicio del Moral, Beatriz; Barzin, Pascal; Beeckman, Bram; BenMoussa, Ali; Berkenbosch, Sophie; Biondi, David; Bonnewijn, Sabrina; Candini, Gian Paolo; Clairquin, Roland; Cubas, Javier; Giordanengo, Boris; Gissot, Samuel; Gomez, Alejandro; Hathi, Brijen; Jeronimo Zafra, Jose; Leese, Mark; Maes, Jeroen; Mazy, Emmanuel; Mazzoli, Alexandra; Meseguer, Jose; Morales, Rafael; Orban, Anne; Pastor-Morales, M; Perez-grande, Isabel; Queirolo, Claudio; Ristic, Bojan; Rodriguez Gomez, Julio; Saggin, Bortolino; Samain, Valérie; Sanz Andres, Angel; Sanz, Rosario; Simar, Juan-Felipe; Thibert, Tanguy

References

- [1] Owen, T., et al. (1988), *Science*, 240, 1767.
- [2] Villanueva, G.L. et al. (2015), *Science* 348, 218.
- [3] Bertaux, J., and F. Montmessin (2001), *J. Geophys. Res.*, 106(E12), 32,879– 32,884.
- [4] Neary, L., and F. Daerden (2018), *Icarus*, 300, 458–476.