

Saturn's Atmospheric Photochemistry: Chemistry and Haze in Ring-Shadowed Atmosphere and within the Hexagon

S.G. Edgington (1), S.K. Atreya (2), E.H. Wilson (3), K.H. Baines (1,4), R.A. West (1), G.L. Bjoraker (5), L.N. Fletcher (6), T. Momary (1)

(1) Jet Propulsion Laboratory/California Institute of Technology, USA (scott.g.edgington@jpl.nasa.gov), (2) University of Michigan, USA, (3) Space Environment Technologies, USA, (4) University of Wisconsin, USA, (5) NASA Goddard Space Flight Center, USA, (6) University of Oxford, UK

Abstract

Cassini has orbited Saturn for over thirteen years. During this Saturn half-year, the ring shadow has moved from covering a relatively large portion of the northern hemisphere (Figure 1a) to covering a large swath of territory south of the equator as solstice approaches. At Saturn Orbit Insertion on July 1, 2004, the sub-solar point was $\sim 24^\circ$ South. At this time, the projection of the optically thick B-ring onto Saturn reached as far as 40° N at the central meridian ($\sim 52^\circ$ N at the terminator). At its maximum extent, the ring shadow can reach as far as 48° N/S ($\sim 58^\circ$ N/S at the terminator). The net result is that the intensity of both ultraviolet and visible sunlight penetrating into any particular latitude will vary greatly depending on both Saturn's axis relative to the Sun and the optical thickness of each ring system. In essence, the rings act like semi-transparent Venetian blinds (Figure 2) over the atmosphere of Saturn.




Figure 2. The optical depth of Saturn's rings in the ultraviolet (Josh Colwell, *pers. comm.*) The rings act like a periodic Venetian blind that will shield atmospheric molecules from solar photons.




Figure 1. Saturn's atmosphere changes in response to the changing inclination of the ring plane relative to the Sun: (a) December 14, 2004, (b) March 16, 2006, (c) April 23, 2008, (d) July 6, 2011, (e) July 29, 2013, (f) November 24, 2014. Images are courtesy of NASA/JPL/Space Science Institute.

Previous work [1,2] examined the variation of the solar flux as a function of solar inclination, i.e. ~ 7.25 year season (Figure 3) at Saturn. Beginning with methane, phosphine and ammonia, we investigate the impact on production and loss rates of the long-lived photochemical products leading to haze formation at several latitudes over a Saturn year. Here, we report on how the oscillating ring shadow modifies the photolysis and production rates of hydrocarbons in Saturn's stratosphere and upper troposphere, including acetylene, ethane, and propane. The magnitude of reflected UV sunlight from the rings on the atmosphere illuminated by ringshine is also explored in this work. Similarly, we assess the impact of these effects on phosphine abundance, a disequilibrium species whose presence in the upper troposphere is a tracer of convection processes in the deep atmosphere. Comparison to the corresponding rates for the clear atmosphere and for the case of

Jupiter, where the solar insolation is known to be insignificant ($\sim 3^\circ$ inclination), will also be presented. Our ongoing analysis of Cassini's CIRS, UVIS, and VIMS datasets that provide abundances of key molecules and estimates of the evolving haze content of the both hemispheres is updated.




Figure 3. This plot illustrates the fraction of Saturn's day that is illuminated by the Sun as a function of solar declination, i.e. season. The curves correspond to sub-solar points of 26.7°S (solid), 19.6°S (dotted), 10.7°S (dashed), and 3.5°S (dot-dashed). Ultimately, this will determine the flux of photons allowed to enter the atmosphere relative to those of a clear, unshaded atmosphere.

The implications for dynamical mixing on the transport of molecules and haze is explored. In particular, we will examine how the now famous hexagonal jet stream (Figure 4) acts like a barrier to transport, isolating Saturn's north polar region from outside transport of photochemically-generated molecules and haze. Future research will explore the role to which increasingly intense sunlight plays in the buildup of hydrocarbon hazes in the polar region and determine exactly how isolated polar region is unaffected by transport from more southerly latitudes.




Figure 4. One of the aims of this exercise is to characterize the haze content Saturn's atmosphere. Several Cassini data sets from ISS (above) and VIMS (below) are being used to meet this goal.

Acknowledgements

The research described in this paper was carried out in part at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration. Copyright 2017, California Institute of Technology. Government sponsorship is acknowledged.

References

- [1] Edgington, S.G., A.A. Simon-Miller, R. Achterberg, G. Bjoraker, P. Romani, F.M. Flasar, J. Colwell, 2006. Adaptation of a 2-D Photochemical Model to Improve Our Understanding of Saturn's Atmosphere. *Bull. American Astron. Soc.*, **38**, 499 (#11.23).
- [2] Edgington, S.G., R.A. West, K.H. Baines, S.K. Atreya, E.H. Wilson, G.L. Bjoraker, L.N. Fletcher, and T. Momary, 2012. Photochemistry in Saturn's Ring Shadowed Atmosphere: Modeling, Observations, and Preliminary Analysis. *Bull. American Astron. Soc.*, **38**, 499 (#11.23).