

Martian aerosol-free reflectance spectra as input to better constrain atmospheric dust content in the NOMAD/TGO nadir observations

Francesca Altieri (1), Valérie Wilquet (2), Fabrizio Oliva (1), Emiliano D'Aversa (1), Filippo Giacomo Carrozzo (1), Giancarlo Bellucci (1), Giuseppe Sindoni (1), Anna Geminale (1), Manish R. Patel (3), Ian R. Thomas (2), Yannick Willame (2), Shohei Aoki (2, 4, 5), Cédric Depiesse (2), Edward Cloutis (2), Frank Daerden (2), Will Hewson (3), Ozgur Karatekin (6), Orietta Lanciano (7), Arnaud Mahieux (2, 5, 8), Jon Mason (3), José Juan Lopez-Moreno (9), Manuel López-Puertas (9), Arianna Piccialli (2), Bojan Ristic (2), Séverine Robert (2), Frédéric Schmidt (10), Ann Carine Vandaele (2), and the NOMAD Team

(1) INAF-IAPS, Rome, Italy, (2) Royal Belgian Institute for Space Aeronomy, BIRA-IASB, Belgium, (3) Open University, United Kingdom, (4) Fonds National de la Recherche Scientifique, Belgium, (5) Tohoku University, Japan, (6) Royal Observatory of Belgium, KSB-ORB, Belgium, (7) ASI, Rome, Italy, (8) University of Texas at Austin, Austin, USA, (9) IAA-CSIC, Spain, (10) Université Paris Sud, Orsay, France (francesca.altieri@iaps.inaf.it)

Abstract

The Martian atmosphere is characterized by the presence of dust particles that during dust storms (global or regional) can reach opacities > 1 . The aim of the present work is to identify and validate a strategy to better constrain atmospheric dust content in the NOMAD/TGO nadir observations making use of aerosol-free reflectance spectra derived from the OMEGA/MEX data set.

1. Rationale

NOMAD, the "Nadir and Occultation for Mars Discovery" experiment [1], is a suite of spectrometers on board the ExoMars Trace Gas Orbiter (TGO) mission that will operate in nadir, limb and solar occultation viewing modes. Nadir spectra will be collected by the Limb, Nadir and Occultation (LNO) and Ultraviolet and Visible Spectrometer (UVIS) channels, covering respectively the 2.3-3.8 μm and the 0.2-0.65 μm spectral ranges. In the visible and infra-red ranges covered by NOMAD dust does not show well-defined absorption bands, although this indication is being tested by other members of the NOMAD team. Airborne dust particles can increase/decrease the albedo for dark/bright terrains and introduce specific slopes depending on opacity and grain size. Surface reflectance in aerosol-free conditions is thus a crucial parameter in the retrieval codes used to assess the dust content in nadir viewing mode. The OMEGA (Observatoire pour la Minéralogie, l'Eau, les Glaces et l'Activité [2]) instrument on board the Mars

Express mission investigated Martian surface properties on a global scale between 0.4-5.1 μm in nadir-pointing geometry and provide a valuable dataset to derive reflectance spectra of the surface in dust-free conditions [e.g. 3].

2. Method

Figure 1 shows two spectra from OMEGA sessions collected in two different seasons but covering the same location (LON= 68.6°E, LAT =15.9°N).




Figure 1: An example of OMEGA spectra, showing the overlap with NOMAD ranges.

The thin spectrum represents the atmosphere corrected reflectance retrieved from the OMEGA session ORB2458_5 using the SAS (Surface

Atmosphere Separation) method [3]. Here, atmospheric features (dust and CO₂ absorptions at 1.5 μm and 2.2 μm) have been removed. The thick spectrum has been measured at the same location during the OMEGA session ORB4558_4 but presents a higher dust content. It represents the top of dusty atmosphere, showing the CO₂ atmospheric features as well. For the dusty spectrum, opacity of the order of 3 has been estimated by means of the method described in [4], using the thin spectrum as input for the surface in dust-free conditions.

3. Summary

Aerosol-free reflectance spectra derived from the OMEGA data set can be used as input for the surface reflectance into NOMAD retrieval codes. Uncertainties due to the different spectral ranges covered by NOMAD and OMEGA will be evaluated and reported in this work. Water ice cloud effects will be also investigated. Improvements in the aerosol properties gathered thanks to NOMAD limb/occultation measurements will be taken into account.

Acknowledgements

This work was supported by Italian Space Agency under grant 2018-2-HH.0. The NOMAD experiment is led by the Royal Belgian Institute for Space Aeronomy (IASB-BIRA), assisted by Co-PI teams from Spain (IAA-CSIC), Italy (INAF-IAPS), and the United Kingdom (Open University). This project acknowledges funding by the Belgian Science Policy Office (BELSPO), with the financial and contractual coordination by the ESA Prodex Office (PEA 4000103401), by Spanish Ministry of Economy, Industry and Competitiveness and by FEDER funds under grant ESP2015-65064-C2-1-P (MINECO/FEDER), as well as by UK Space Agency through grant ST/P000886/1. This work was supported by the European Union's Horizon 2020 Programme under grant agreement UPWARDS-633127.

The NOMAD Team

Scientific team: Vandaele, Ann Carine; Lopez Moreno, Jose Juan; Bellucci, Giancarlo; Patel, Manish; Allen, Mark; Alonso-Rodrigo, Gustavo; Altieri, Francesca; Aoki, Shohei; Bauduin, Sophie; Bolsée, David; F. Giacomo Carrozzo, Clancy, Todd; Cloutis, Edward; Daerden, Frank; D'Aversa,

Emiliano; Depiesse, Cédric; Erwin, Justin; Fedorova, Anna; Formisano, Vittorio; Funke, Bernd; Fussen, Didier; Garcia-Comas, Maia; Geminale, Anna; Gérard, Jean-Claude; Gillotay, Didier; Giuranna, Marco; Gonzalez-Galindo, Francisco; Hewson, Will; Homes, James; Ignatiev, Nicolai; Kaminski, Jacek; Karatekin, Ozgur; Kasaba, Yasumasa; Lanciano, Orietta; Lefèvre, Franck; Lewis, Stephen; López-Puertas, Manuel; López-Valverde, Miguel; Mahieux, Arnaud; Mason, Jon; Mc Connell, Jack; Mumma, Michael J.; Nakagawa, Hiromu, Neary, Lori; Neefs, Eddy; Novak, R.; Oliva, Fabrizio; Piccialli, Arianna; Renotte, Etienne; Robert, Severine; Sindoni, Giuseppe; Smith, Mike; Stiepen, Arnaud; Thomas, Ian; Trokhimovskiy, Alexander; Vander Auwera, Jean; Villanueva, Geronimo; Viscardy, Sébastien; Whiteway, Jim; Willame, Yannick; Wilquet, Valérie; Wolff, Michael; Wolkenberg, Paulina – *Tech team:* Alonso-Rodrigo, Gustavo; Aparicio del Moral, Beatriz; Barzin, Pascal; Beeckman, Bram; BenMoussa, Ali; Berkenbosch, Sophie; Biondi, David; Bonnewijn, Sabrina; Candini, Gian Paolo; Clairquin, Roland; Cubas, Javier; Giordanengo, Boris; Gissot, Samuel; Gomez, Alejandro; Hathi, Brijen; Jeronimo Zafra, Jose; Leese, Mark; Maes, Jeroen; Mazy, Emmanuel; Mazzoli, Alexandra; Meseguer, Jose; Morales, Rafael; ; Orban, Anne; Pastor-Morales, M; Perez-grande, Isabel; Queirolo, Claudio; Ristic, Bojan; Rodriguez Gomez, Julio; Saggin, Bortolino; Samain, Valérie; Sanz Andres, Angel; Sanz, Rosario; Simar, Juan-Felipe; Thibert, Tanguy

References

- [1] Vandaele, A.C., Neefs, E., Drummond, R. et al.: Science objectives and performances of NOMAD, a spectrometer suite for the Exo-Mars TGO mission, *Planetary and Space Science*, 119, 233–249, 2015.
- [2] Bibring, J-P., et al.: Omega: Observatoire pour La minéralogie, l'eau, Les Glaces Et l'activité. ESA SP-1240: Mars Express: The Scientific Payload. ESA Publications Division, ESTEC, Noordwijk, The Netherlands, pp. 37–49, 2004.
- [3] Geminale, A., et al.: Removal of atmospheric features in near infrared spectra by means of principal component analysis and target transformation on Mars: I method. *Icarus* 253, 51–65, 2015.
- [4] Oliva, F., et al.: Properties of a Martian local dust storm in Atlantis Chaos from OMEGA/MEX data, *Icarus*, 300, 1–11, 2018.