



Patterns in textured dust storms in Mars North Pole

Agustin Sanchez-Lavega¹, Josu Garcia-Morales², Jorge Hernandez-Bernal^{1,2}, Tera del Rio-Gaztelurrutia¹, Ricardo Hueso¹, Eleni Ravanis³, Alejandro Cardesin-Moinelo³, Dimitri Titov⁴, Simon Wood⁵, Daniela Tirsch⁶, Ernst Hauber⁶, and Klaus-Dieter Matz⁶

¹Universidad Pais Vasco UPV/EHU, Escuela de Ingeniería de Bilbao, Física Aplicada I, Bilbao, Spain (agustin.sanchez@ehu.es)

²Aula Espazio Gela, Escuela de Ingeniería de Bilbao, Universidad del País Vasco UPV/EHU, Bilbao, Spain

³European Space Agency, ESAC, Madrid, Spain

⁴European Space Agency, ESTEC, Noordwijk, Netherland

⁵European Space Agency, ESOC, Darmstadt, Germany

⁶German Aerospace Center (DLR), Institute of Planetary Research, Berlin, Germany

We report on the cloud top morphology, scale-analysis of patterns, and dynamics of “textured” local dust storms on Mars observed at the edge of the North Polar cap during the Northern Hemisphere Spring Equinox, before aphelion, using images obtained by the Visual Monitoring Camera (VMC) [1] and High Resolution Stereo Camera (HRSC) [2] onboard Mars Express. VMC images were analyzed with tools described in previous works [3-4] and HRSC images were analyzed from map-projections.

The observations cover the period from March 3 to July 17, 2019, corresponding to the solar longitude range $L_s = 350^\circ - 55^\circ$ (Martian Years 34 to 35). We observed the continuous formation of circumpolar dust patches, large frontal arc-shaped features, flushing dust storms, textured local dust storms and other forms of cloud activity at the edge of and inside the North Polar cap around latitude 60°N , a rich phenomenology typical of this season [5]. In this presentation we concentrate on the study of three textured local dust storms observed at the end of May and early June 2019.

The observed textured storms contained cellular structure and frontal-like banding, both indicative of organized active lifting within the storm [6-7]. The first storm was centered at about 185°E , 60°N and occupied a small area of $1.75 \times 10^5 \text{ km}^2$. It showed three frontal bands with lengths $\sim 1000 \text{ km}$ and widths of 85 km separated by 40 km . In the interior of the storm a cellular pattern developed with a mean size of $50 \text{ km} \times 20 \text{ km}$. The second storm was centered at about 330°E , 60°N , occupied an area of $1.3 \times 10^6 \text{ km}^2$ and moved zonally with velocities from 20 to 45 ms^{-1} . A global cellular pattern covered this storm with cells of a mean size of $135 \text{ km} \times 60 \text{ km}$ and inter-cell distances in the range $100\text{-}300 \text{ km}$. The third storm was centered at about 150°E , 65°N , occupied an area of $1.6\text{-}2.1 \times 10^5 \text{ km}^2$ and moved zonally with a mean velocity of 38 ms^{-1} . Its cellular pattern had a mean size of $70 \text{ km} \times 40 \text{ km}$. In all cases, the cell texture is anisotropic in the horizontal size (length/width, $l/w \sim 2$) and their value is well above the atmospheric scale height ($H \sim 8 \text{ km}$). Deep convection driven by buoyancy generated by the radiative heating of atmospheric dust is proposed to explain this structure.

References:

[1] Ormston, T., Denis, M., Scuka, D., & Griebel, H., An ordinary camera in an extraordinary location: Outreach with the Mars Webcam. *Acta Astronautica*, **69**, 703-713 (2011)

[2] Jaumann, R., Neukum, G., Behnke, T., Duxbury, T. C., Eichtenopf, K., Flohrer, et al., The high-resolution stereo camera (HRSC) experiment on Mars Express: Instrument aspects and experiment conduct from interplanetary cruise through the nominal mission. *Planetary and Space Science*, **55**, 928-952 (2007)

[3] Sánchez-Lavega, A., Chen-Chen, H., Ordonez-Etxeberria, I., Hueso, R., del Río-Gaztelurrutia, T., Garro, A., & Wood, S., Limb clouds and dust on Mars from images obtained by the Visual Monitoring Camera (VMC) onboard Mars Express. *Icarus*, **299**, 194-205 (2018)

[4] Hernández-Bernal, J., Sánchez-Lavega, A., del Río-Gaztelurrutia, T., Hueso, R., Cardesín-Moinelo, A., Ravanis, E. M., Tivov, D., & Wood, S., The 2018 Martian Global Dust Storm over the South Polar Region studied with MEx/VMC. *Geophys. Research Lett.*, **46**, 10330-10337 (2019)

[5] Kahre, M. A., Murphy, J. R., Newman, C. E., Wilson, R. J., Cantor, B. A., Lemmon, M. T. & Wolff, M. J., The Mars Dust Cycle. In R. Haberle, R. T. Clancy, F. Forget, M. D. Smith and R. W. Zurek (Eds.), *The Atmosphere and Climate of Mars* (pp. 295-337). Cambridge, U.K. Cambridge University Press. (2017)

[6] Guzewich S. D., Toigo A. D., Kulowski L., Wang H., Mars Orbiter Camera climatology of textured dust storms, *Icarus*, **258**, 1-13 (2018)

[7] Heavens N. G., Textured Dust Storm Activity in Northeast Amazonis–Southwest Arcadia, Mars: Phenomenology and Dynamical Interpretation, *J. Atmos. Sci.*, **74**, 1011-1037 (2017)