



## **Reflected, transmitted irradiance, radiative divergence and heating rate on the base of simple atmospheric optical models**

I. Melnikova (1) and A. Ginzburg (2)

(1) Russian Academy of Sciences, Research Center for Interdisciplinary Environmental Cooperation (INENCO RAS), Laboratory for Atmospheric Physics, Sankt-Petersburg, Russian Federation (irina.melnikova@pobox.spbu.ru), (2) Russian Academy of Sciences, A.M.Obukhov Institute of Atmospheric Physics, Moscow, Russia (gin@ifaran.ru)

Rapid evolution of computers in last decades provides complicating atmospheric models with detailing vertical profiles, accounting for irregular clouds in wide spectral ranges. Numerical algorithms for calculating radiative characteristics with maximal exactness and minimization of uncertainty are usually applied. There are many different computer codes including look-up tables with aerosols characteristics, water vapor, atmosphere conditions in different latitudes, continental and sea conditions, and seasons. Sophisticated approaches for calculating optical parameters are based on scattering and radiative transfer theories. It is very useful for applied problems. However the analysis of separate factors influence on atmospheric radiative characteristics without of considering all possible variations of the whole totality is often necessary for many research problems. For that case the simple models of homogeneous (for the clear atmosphere) and two or three layer atmosphere (for cloudy cases) allow operative varying considered atmospheric optical parameters and provide result that hardly yields to complicate models and clearly elucidating interactions between of key atmospheric parameters and radiative characteristics. Two-stream methods of radiative transfer theory ensure an acceptable exactness for calculating integral (over viewing directions) radiative characteristics (irradiance and radiative divergences). Asymptotic formulas are effective for fast and transparent calculation in case of the cloud atmosphere.

Hence we accept a simplest optical model of the homogeneous clear atmosphere including ozone absorption in UV ranges, molecular scattering, and two variants of the aerosol content at selected shortwave wavelength. In cloud case three variants of extended cloud layer are added. Radiative characteristics together with heating rate are calculated and presented in this study.