



Analyses of possible changes in the mean and extreme precipitation regimes over Spain under climate change scenarios.

M. Turco (1), M.C. Llasat (1), and P. Quintana Seguí (2)

(1) University of Barcelona, Department of Astronomy and Meteorology, Barcelona, Spain (mturco@am.ub.es; +34-(0)93-4021133), (2) Observatori de l'Ebre (Universitat Ramon Llull - CSIC), Roquetes, Spain.

Global climate models (GCM) are basic tools to study the climate. However due to their coarse resolution, generally few hundred kilometres (Solomon et al. 2007), they are not useful to study regional aspects of the climate of order of tens of kilometres (Cohen, 1990). This is especially true for Spain, which is geographically complex and heterogeneous and it is characterized by a great variability of precipitation (Serrano et al. 1999, Trigo and Palutikof, 2001). Consequently, developing regional climate scenarios is a key aspect for any impact and adaptation study to climate change in Spain.

In contrary to the traditional approach in which the statistical downscaling has been considered as an alternative to dynamical downscaling, we used the two kinds of downscaling approaches together in order to combine their advantages. This hybrid approach is becoming popular (Maraun et al. 2010, Piani et al. 2010, Quintana Seguí et al 2010, Themeßl et al. 2010) thank to the better skill of RCMs (e.g. Herrera et al. 2010a) and the increase of availability of RCMs (e.g. due to projects like ENSEMBLES).

In this study, the fields of precipitation simulated by an ensemble of RCMs are statistically post-processed in order to downscale and to calibrate their precipitation fields using the analog method, (Lorenz, 1969), according to the approach of the Model Output Statistics (MOS, Wilks, 2006). The simulations used as predictors have been the RCMs provided by the EU-funded project ENSEMBLES. The Spain02 precipitation dataset (Herrera et al. 2010b) is used as predictand. The application of the MOS-analog method shows a big reduction of the systematic error of the RCMs in representing the spatial pattern of the climatology of several standard and extreme precipitation indices (Turco et al. 2011, accepted in JGR).

After an extensive validation of this method under optimal (using RCMs driven by ERA40) and sub-optimal conditions (using RCMs driven by GCM in current climate), we have also tested the validity of the statistical relationship in a surrogate climate as in (Frias et al. 2006). Finally, we present the application of this method to RCM scenario run.