

Ozone time scale decomposition and trend assessment from surface observations in Switzerland and Europe

E. Boleti (1), C. Hüglin (1), and S. Takahama (2)

(1) Empa, Swiss Federal Laboratories for Materials Science and Technology, Dübendorf, Switzerland (eirini.boleti@empa.ch, Christophe.Hueglin@empa.ch), (2) EPFL, Ecole Polytechnique Federale de Lausanne, Switzerland (satoshi.takahama@epfl.ch)

Emissions of ozone precursors have been regulated in Europe since around 1990 with control measures primarily targeting to industries and traffic. In order to understand how these measures have affected air quality, it is now important to investigate concentrations of tropospheric ozone in different types of environments, i.e. urban, suburban and rural and in different geographic regions. In this study, we analyse high quality data sets for Switzerland (NABEL network) and whole Europe (AirBase) for the last 25 years to calculate long-term trends of ozone concentrations. A sophisticated time scale decomposition method, called the Ensemble Empirical Mode Decomposition (EEMD) [Huang et al. (1998), Wu and Huang (2009)], is used for decomposition of the different time scales of the variation of ozone, namely the long-term trend, seasonal and short-term variability. This allows subtraction of the seasonal pattern of ozone from the observations and estimation of long-term changes of ozone concentrations. We observe that, despite the implementation of regulations, for most of the measurement sites ozone daily mean values have been increasing almost until mid-2000s. Afterwards, we observe a decline or a leveling off in the concentrations; certainly a late effect of limitations in ozone precursor emissions. On the other hand, the peak ozone concentrations have been decreasing for almost all regions. The evolution in this trend exhibits some differences between the different types of measurement locations as well as geographic regions. In addition, ozone is known to be strongly affected by meteorology. In the applied approach, some of the meteorological effects are already captured by the seasonal signal and already removed in the de-seasonalized ozone time series. For adjustment of the influence of meteorology on the higher frequency ozone variation, a statistical approach, which corrects for meteorological effects, has been developed in order to investigate a) if trends are masked by meteorological variability and b) to understand which part of the observed trends is meteorology driven. Generalized Additive Models (GAM)[Hastie and Tibshirani (1990), Wood (2006)] perform very well in accounting the relationship between air pollutants' concentrations and meteorological parameters. In our case, by correlating short-term variation of ozone, as obtained from the EEMD, with the corresponding short-term variation of relevant meteorological parameters, we subtract the variation of ozone concentrations that is related to the meteorological effects explained by the GAM. We find that higher frequency meteorological correction does not have a large effect on the trends of the daily mean concentrations, and the variance in the meteorology corrected time series is only reduced by a small factor. The influence of the correction is slightly larger for the maximum values of ozone. In conclusion, we present a sophisticated and consistent approach for detecting and categorizing trends and meteorological influences on ozone concentrations in long-term measurements across Europe.

References

[Hastie and Tibshirani, 1990] Hastie, T. and Tibshirani, R. (1990). Generalized Additive Models. *Statistical Science*, 1(3):297–310.

[Huang et al., 1998] Huang, N. E., Shen, Z., Long, S. R., Wu, M. C., Shih, H. H., Zheng, Q., Yen, N. C., Tung, C. C., and Liu, H. H. (1998). The empirical mode decomposition and the Hilbert spectrum for nonlinear and non-stationary time series analysis. *Proceedings of the Royal Society of London. Series A: Mathematical, Physical and Engineering Sciences*, 454(1971):903–995.

[Wood, 2006] Wood, S. (2006). *Generalized Additive Models: An Introduction with R*. Chapman & Hall/CRC.

[Wu and Huang, 2009] Wu, Z. and Huang, N. E. (2009). Ensemble Empirical Mode Decomposition. *Advances in Adaptive Data Analysis*, 1(1):1–41.