

## Revisiting the Schönbein ozone measurement methodology

I. A. Ramírez-González (1), J. A. Añel (1,2), A. Saiz-López (3), O. Garcia-Feal (1), A. Cid (4,5), J. C. Mejuto (5), and L. Gimeno (1)

(1) EPhysLab, Fac. of Sciences of Ourense, Universidade de Vigo, Ourense, Spain, (2) Smith School of Enterprise and the Environment, University of Oxford, Oxford, United Kingdom, (3) Department of Atmospheric Chemistry and Climate, Institute of Physical Chemistry Rocasolano, CSIC, Madrid, Spain, (4) UCIBIO, REQUIMTE, Departamento de Química, Faculdade de Ciências e Tecnologia, Universidade NOVA de Lisboa, Caparica, Portugal., (5) Department of Physical–Chemistry, Fac. of Sciences of Ourense, Universidade de Vigo, Ourense, Spain

Trough the XIX century the Schönbein method gained a lot of popularity by its easy way to measure tropospheric ozone. Traditionally it has been considered that Schönbein measurements are not accurate enough to be useful. Detractors of this method argue that it is sensitive to meteorological conditions, being the most important the influence of relative humidity. As a consequence the data obtained by this method have usually been discarded. Here we revisit this method having into account that values measured during the 19th century were taken using different measurement papers. We explore several concentrations of starch and potassium iodide, the basis for this measurement method. Our results are compared with the previous ones existing in the literature. The validity of the Schönbein methodology is discussed having into account humidity and other meteorological variables.